

Partial Identification in Monotone Binary Models: Discrete Regressors and Interval Data.

Thierry Magnac* Eric Maurin†

First version: February 2004

This revision: October 2005

Abstract

We investigate identification in semi-parametric binary regression models, $y = 1(x\beta + v + \epsilon > 0)$ when ϵ is assumed uncorrelated with a set of instruments z , ϵ is independent of v conditionally on x and z , and the conditional support of ϵ is sufficiently small relative to the support of v . We characterize the set of observationally equivalent parameters β when v is discrete or when interval data only are available on v . When there exist as many instruments z as variables x , the sets within which lie the scalar components β_k of parameter β can be estimated by simple linear regressions. Also, in the case of interval data, it is shown that additional information on the distribution of v within intervals shrinks the identification set. Namely, the closer to uniformity the distribution of v is, the smaller the identification set is. Point identification is achieved if and only if v is uniform within intervals.

*University of Toulouse (GREMAQ and IDEI), 21 Allée de Brienne, 31000 Toulouse, France, *Address:* magnac@cict.fr, <http://www.idei.fr/vitae/magnac.html>

†PSE, 48 Bld Jourdan, 75014, Paris, France, *Address:* eric.maurin@ens.fr

1 Introduction¹

In empirical research, point identification of parameters often requires assumptions that are difficult to motivate. Yet, the recourse to more credible and weaker restrictions that lead to partial identification remains rare although the context of partial identification, when the identification region is convex and bounded, is not conceptually different from the familiar context of confidence intervals.

We analyse in this paper partial identification in a binary regression model with discrete or interval-valued data, a case that is admittedly specific although surprisingly rich in terms of implications and facility of application. The identification set is bounded and convex and bounds are easy to characterise by simple moments of the data.

One of the practical interest of our model stems from the fact that the data on covariates that researchers have access to, are very often discrete or interval-valued. Such covariates tend to render point identification very problematic. When all covariates are discrete or when only interval data are available, point identification of parameters of popular index models is lost whatever the identifying restrictions (Manski (1988)). When all covariates (denoted x) are discrete, Bierens and Hartog (1988) have indeed shown that there exists an infinite number of single-index representations for the mean regression of a dependent variable, y , i.e. $E(y | x) = \varphi_\theta(x\theta)$. Under weak conditions, the set of observationally equivalent parameters θ is actually dense in its domain of variation, Θ . Put differently, for almost any parameter θ there exists a measurable real function φ_θ such that $E(y | x)$ can be written $\varphi_\theta(x\theta)$.

A recent contribution by Manski and Tamer (2002) considers a reasonably more specific framework where the non-parametric mean regression $E(y | x)$ is assumed monotonic with respect to at least one particular regressor, say v . As a particular case, they study inference on the parameter of the familiar semiparametric binary regression model $y = 1(x\beta + v + \epsilon > 0)$, when v is not perfectly observed, i.e., when interval-data only are available on v . They analyse the identification of β under the quantile independence assumption underlying the analysis of Manski (1985). Most interestingly, they show that this assumption restricts the model parameters to a non-empty, convex set of observationally equivalent values that they

¹We thank Arthur Lewbel for helpful discussions and participants at seminars at LSE, CREST, CEMFI, Montréal and Rochester and at conferences (ESRC Econometrics Study Group in Bristol, ESWC'05) for helpful comments. The usual disclaimer applies.

characterise. Yet, the estimation of the identification region is not straightforward. It requires a modification of the maximum score technique introduced by Manski (1985).

The first message of our paper is that set-identification of parameter β in the binary model can be obtained through a different set of restrictions (i.e. Lewbel, 2000), and that the estimation of the identification region in this setting only requires usual regression tools. Specifically, it is shown that the combination of an uncorrelated-error assumption (i.e., $E(x'\varepsilon) = 0$) with a partial independence assumption (i.e., $F_\varepsilon(\varepsilon | x, v) = F_\varepsilon(\varepsilon | x)$) and a support assumption ($Supp(-x\beta - \varepsilon) \subset [v_l, v_u]$, where v_l and v_u are finite) restricts the parameters of the semiparametric binary regression model to a non-empty bounded and convex set. We characterise this set and show that the bounds of the intervals in which lie any scalar linear combination of parameter β can be estimated through simple linear regression methods. These findings are probably the most important results of this paper for practitioners because they enable researchers to estimate the identified set very easily.

Partial independence implies that the binary outcome is monotone in v . Interestingly, the support condition does not generate specific restrictions on the binary data that can be analysed although it requires to be careful when it comes to application. The support assumption implies that the probability of success $\Pr(y = 1 | x, v)$ increases from 0 and 1 when v varies between the lower and upper values of v (v_l and v_u) although those values are not necessarily observed in the data. Potential applications of this set-up include the analysis of the probability of buying an object as a function of price, discrete by construction, or the analysis of the diffusion of a binary outcome as a function of secular time-trend or the study of entry into (or exit from) the school system or the labor market as a function of age. For example, consider the analysis of the school-leaving probability, where v stands for age. For institutional reasons, we may posit that this probability varies from zero to one when v varies from 16 years to 30 years so that parameter β is partially identified. The width and location of the identification region will depend on how age is measured including the range of variation of individual ages actually observed in the sample.

The last interesting feature of the set-up analysed in this paper is that, in the case where v is censored by interval, additional information on the distribution of v within intervals might reduce the size of the identification region. Specifically, the size of the identification

region diminishes, in a sense made precise below, as the distribution of the special regressor within intervals becomes closer to uniformity. It is shown that the identification set is a singleton and the parameter of interest β is exactly identified if and only if v is uniformly distributed within intervals. This property is particularly interesting when one has control over the process of censoring the continuous data on v (*e.g.* the birthdate) into interval data (*e.g.* month of birth). In order to minimize the size of the identification set, one should censor the data in such a way that the distribution of the censored variable is as close as possible to a uniform distribution within the resulting intervals.

As for references, this paper belongs to the small, but growing literature on partial identification as pioneered by Manski (2003, and references therein). Our results on bounds on parameters in binary regressions can be seen as generalizations of bounds on averages, derived in the paper by Green, Jacowitz, Kahneman & McFadden (1998). They are also reminiscent of the results presented by Leamer (1987) in a system of equations where covariates are mismeasured. There, the vector of parameters of interest lies in an ellipsoid though it is not known whether intervals obtained by projecting this ellipsoid onto individual directions can be estimated directly. Also, there exist striking similarities between our identification results and those of Chesher (2005), even though the topic is quite different. Chesher estimates the local effect of an endogenous discrete variable in non-separable models and shows that discrete variation of this endogenous variable as opposed to continuous variation is likely to give rise to partial identification.

Recent applications of the partial independence setting include Lewbel (2003) who studies the probability of obtaining a university degree using the cost of attending a local public college (relative to local unskilled wages) as exogenous regressor v or Maurin (2002) who estimates an education production function using the date of birth within the year as exogenous regressor. In contingent valuation studies, where ε stands for unobserved taste variation, v can be the bid that is determined by experimental design, and so may be constructed by the researcher to satisfy the necessary exclusion and support restrictions. Lewbel, Linton and McFadden (2001) provide an empirical application for this case.

This paper focuses on identification issues, not on inference problems. As a matter of fact, the inference issues are addressed by recent works by Chernozhukov, Hong and Tamer (2004),

Imbens and Manski (2004) or Horowitz and Manski (2000). In particular, Chernozhukov et al. (2004) study inference in multivariate cases under more general conditions of set-identification than ours and their findings can be applied to our results. It is also possible to follow Horowitz and Manski (2000) who study inference about intervals through inference on the lower and upper bounds of identified intervals. An alternative route is proposed by Imbens and Manski (2004) who changed focus by considering inference about the true value of the parameter (within an identified interval) and not about the interval in itself.

The paper is organized as follows : The first section sets up notations and models. The second section examines the discrete case, the third section analyzes the case of interval data, the fourth section reports Monte Carlo experiments and the last section concludes. Since the abstract case where the x are endogenous is not more complex than the case where they are exogenous, we will consider right from the start the endogenous case where ϵ , though potentially correlated with the variables x , is uncorrelated with a set of instruments z . Some results will be specialized to the case where the number of instruments is equal to the number of explanatory variables. All proofs are in appendices.

2 The Set-Up

Let the “data” be given by the distribution of the following random variable²:

$$\omega = (y, v, x, z)$$

where y is a binary outcome, while v , x and z are covariates and instrumental variables which role and properties are specified below. We first introduce some regularity conditions on the distribution of ω . They will be assumed valid in the rest of the text.

Assumption R(egularity):

R.i. (Binary model) The support of the distribution of y is $\{0, 1\}$

R.ii. (Covariates & Instruments) The support of the distribution, $F_{x,z}$ of (x, z) is $S_{x,z} \subset \mathbb{R}^p \times \mathbb{R}^q$. The dimension of the set $S_{x,z}$ is $r \leq p + q$ where $p + q - r$ are the potential overlaps and functional dependencies.³ The condition of full rank, $rank(E(z'x)) = p$, holds.

²We only consider random samples and we do not subscript individual observations by i .

³With no loss of generality, the p explanatory variables x can partially overlap with the $q \geq p$ instrumental

R.iii. (Discrete or Interval-Valued Regressor) The support of the distribution of v conditional on (x, z) is $\Omega_v \subset \mathbb{R}$ almost everywhere- $F_{x,z}$ (a.e. $F_{x,z}$). This conditional distribution, denoted $F_v(\cdot | x, z)$, is defined a.e. $F_{x,z}$. In the remainder we will assume either $\Omega_v = \{v_1, \dots, v_K\}$ (discrete case) or $\Omega_v = [v_1, v_K[$ (interval case) where v_1 and v_K are finite. In both cases Ω_v^0 will denote $[v_1, v_K[$.

R.iv. (Functional Independence) There is no subspace of $\Omega_v \times S_{x,z}$ of dimension strictly less than $r + 1$ which probability measure, $(F_v(\cdot | x, z).F_{x,z})$, is equal to 1.

Assumption *R.i* defines a binary model where there are p explanatory variables and q instrumental variables (assumption *R.ii*). Given assumption *R.ii*, we could denote the functionally independent description of (x, z) as u and this notation could be used interchangeably with (x, z) . In assumption *R.iii*, the support of v is assumed to be independent of variables (x, z) . If this support is an interval in \mathbb{R} (including \mathbb{R} itself) and v is continuously observed, we are back to the case studied by Lewbel (2000) and Magnac & Maurin (2004). In the next section (section 3), this support is assumed to be discrete so that the special regressor is said to be discrete. In section 4, the support is assumed continuous, but v is observed imperfectly because it is censored. In such a case, the special regressor is said to be interval-valued. In all cases, Assumption *R.iv* avoids the degenerate case where v and (x, z) are functionally dependent.

There are many examples of discrete covariates in applied econometrics. Variables such as gender, levels of education, occupational status or household size of survey respondents typically take a discrete number of values. In contingent valuation studies, prices are set by the experimenter and they are in general discrete, by steps of 0.10, 1 or 10 euros. There are also many examples of interval-valued data. They are common in surveys where, in case of non-response to an item, follow-up questions are asked. Manski & Tamer (2002) describe the example of the Health and Retirement Study. If a respondent does not want to reveal his wealth, he is then asked whether it falls in a sequence of intervals (“unfolding brackets”). Another important reason for interval data is anonymity. Age is a continuous covariate which could in theory be used as a source of continuous exogenous variation in

variables z . Variables (x, z) may also be functionally dependent (for instance $x, x^2, \log(x), \dots$). A collection (x_1, \dots, x_K) of real random variables is functionally independent if its support is of dimension K (i.e. there is no set of dimension strictly lower than K which probability measure is equal to 1).

many settings. For confidentiality reasons however, the French National Statistical Office censors this information in the public versions of household surveys by transforming dates of birth into months (or years) of birth only. French statisticians are afraid that the exact date of birth along with other individual and household characteristics might reveal the identity of households responding to the survey.

Assuming that the data satisfy *R.i–R.iv*, the basic issue addressed in this paper is whether they can be generated by the following semi-parametric latent variable index structure :

$$y = \mathbf{1}\{x\beta + v + \epsilon > 0\}, \quad (\text{LV})$$

where $\mathbf{1}\{A\}$ is the indicator function that equals one if A is true and zero otherwise and where the random shock ϵ satisfies the following properties,

Assumption L(atent)

(L.1) (*Partial independence*) The conditional distribution of ϵ given covariates x and variables z is independent of the covariate v :

$$F_\epsilon(\cdot \mid v, x, z) = F_\epsilon(\cdot \mid x, z)$$

The support of ϵ is denoted $\Omega_\epsilon(x, z)$.

(L.2) (*Large support*) The support of $-x\beta - \epsilon$ is a subset of Ω_v^0 as defined in *R(iii)*.

(L.3) (*Moment condition*) The random shock ϵ is uncorrelated with variables z : $E(z'\epsilon) = 0$.

This combination of identifying assumptions was introduced by Lewbel (2000) and Honoré and Lewbel (2002).⁴ Powell (1994) discusses partial independence assumptions (calling them exclusion restrictions) in the context of other semiparametric models, i.e. without combining them with (L.2) or (L.3).

Generally speaking, the relationship between the set-up used in this paper and the assumptions used by Manski and Tamer (2002) are similar to the relationship between Manski

⁴There is only a minor difference between assumption *L* and the set-up introduced by Lewbel (2000), namely the distribution function F_ϵ can have mass points. When the special regressor is discrete or interval-valued, it is much easier than in the continuous case to allow for such discrete distributions of the unobserved factor. If all distribution functions are CADLAG (i.e., continuous on the right, limits on left), the large support assumption (L.2) has to be slightly rephrased however in order to exclude a mass point at $-x\beta - v_K$.

(1985) and Lewbel (2000). The quantile independence set-up assumes that one quantile of ε is independent of all covariates, whereas the partial independence assumption used in this paper is equivalent to assuming that all quantiles of ε are independent of one covariate. In this crude sense, both assumptions are comparably restrictive. Another difference is that the partial independence hypothesis makes it possible to characterize the domain of observationally equivalent distribution functions of the unobserved residuals. The price to pay is that partial independence requires conditions on the support of the covariates that are stronger than the conditions required under quantile-independence. Assumption L and some examples are commented in Lewbel (2000) or Magnac and Maurin (2004). Once v is perfectly observed and continuously distributed, the latter paper shows that Assumption L is sufficient for exact point-identification of both β and $F_\varepsilon(\cdot | x, z)$.

The rest of the paper focuses on the identification of β . Before moving on to this issue, it is important to understand the relationship between this parameter and the effect of changes in the covariates on the choice probability. Let us consider the effect on y of assigning (say) $(x' = x + \Delta x, v, z, \varepsilon)$ to everyone of type (x, v, z, ε) . Using model (LV), the partial independence assumption and integrating out ε , the average effect simply boils down to $E(y | x, v + \beta\Delta x, z) - E(y | x, v, z)$, namely to the expected difference in the conditional probability of success observed in the data between individuals of type $(x, v + \beta\Delta x, z)$ and individuals of type (x, v, z) . Hence, β defines the shifts in v which enable the evaluation of the average effects on y of *ceteris paribus* shifts in x , where the term *ceteris paribus* means that we hold ε and v constant.

In the remainder, any $(\beta, F_\varepsilon(\cdot | x, z))$ satisfying Assumption L is called a latent model. The index parameter $\beta \in \mathbb{R}^p$ is the unknown parameter of interest. The distribution function of the error term, ε , is also unknown and may be considered as a nuisance parameter. Identification is studied in the set of all such $(\beta, F_\varepsilon(\cdot | x, z))$.

3 The Discrete Case

In this section, the support of the special regressor is supposed to be a discrete set given by:

Assumption D(iscrete): $\Omega_v = \{v_1, \dots, v_K\}, v_k < v_{k+1}$ for any $k = 1, \dots, K - 1$.

To begin with, we are going to explore the properties that a conditional probability distribution $Pr(y = 1 | v, x, z)$ necessarily verifies when it is generated by a latent model $(\beta, F_\varepsilon(\cdot | x, z))$ that satisfies Assumption L . The issue is to make explicit the class of binary outcomes which can actually be analyzed through the latent models under consideration.

3.1 Characterizing the Conditional Distribution

As defined by (L.1) and (L.2), partial-independence and large-support assumptions restrict the class of binary outcomes that can actually be analyzed. Restrictions are characterized by the following lemma that is proved in Magnac and Maurin (2004) and that we repeat here for completeness:

Lemma 1 *Under partial independence (L.1) and large support (L.2) conditions, we necessarily have:*

(NP.1) *(Monotonicity) The conditional probability $Pr(y_i = 1 | v, x, z)$ is non decreasing in v (a.e. $F_{x,z}$).*

(NP.2) *(Support) The conditional probability $Pr(y_i = 1 | v, x, z)$ varies from 0 to 1 when v varies over its support:*

$$Pr(y_i = 1 | v = v_1, x, z) = 0, \quad Pr(y_i = 1 | v = v_K, x, z) = 1.$$

If a binary outcome does not satisfy (NP.1) or (NP.2) then there exists no latent model generating the reduced form $Pr(y = 1 | v, x, z)$. The next section studies whether the reciprocal holds true, i.e. whether (NP.1) and (NP.2) are sufficient conditions for identification.

3.2 Set-identification

We consider a binary reduced-form $Pr(y = 1 | v, x, z)$ satisfying the monotonicity condition (NP1) and the support condition (NP2) and ask whether there exists a latent model $(\beta, F_\varepsilon(\cdot | x, z))$ generating this reduced-form through the latent variable transformation (LV). The answer is positive though the admissible latent model is not unique. There are many possible latent models which parameters are observationally equivalent.

We begin with a one-to-one change in variables which will allow us to characterize the set of observationally equivalent parameters through simple linear moment conditions. Denote,

for $k \in \{2, \dots, K - 1\}$:

$$\begin{aligned}\delta_k &= (v_{k+1} - v_{k-1})/2 \\ p_k(x, z) &= Pr(v = v_k \mid x, z).\end{aligned}$$

Using these notations, the transformation of the binary response variable which is adapted to the discrete case, is defined as:⁵

$$\begin{aligned}\tilde{y} &= \frac{\delta_k \cdot y}{p_k(x, z)} - \frac{v_K + v_{K-1}}{2} \text{ if } v = v_k, \text{ for } k \in \{2, \dots, K - 1\}, \\ \tilde{y} &= -\frac{v_K + v_{K-1}}{2} \text{ if } v = v_1 \text{ or } v = v_K,\end{aligned}\tag{1}$$

In contrast to the continuous case, the identification of β when v is discrete is not exact anymore as stated in the following theorem:

Theorem 2 Consider β a vector of parameter and $Pr(y = 1 \mid v = v_k, x, z)$, (denoted $G_k(x, z)$) a conditional probability distribution satisfying conditions of monotonicity (NP.1) and support (NP.2). The two following statements are equivalent,

(i) there exists a latent random variable ε such that the latent model $(\beta, F_\varepsilon(\cdot \mid x, z))$ satisfies Assumption L and such that $\{G_k(x, z)\}_{k=1, \dots, K}$ is its image through the transformation (LV),

(ii) there exists a measurable function $u(x, z)$ from $S_{x,z}$ to \mathbb{R} which takes its values in the interval (a.e. $F_{x,z}$)

$$I(x, z) =] - \Delta(x, z), \Delta(x, z)[,$$

where

$$\Delta(x, z) = \frac{1}{2} \sum_{k=2}^K [(v_k - v_{k-1})(G_k(x, z) - G_{k-1}(x, z))],$$

and such that,

$$E(z'(x\beta - \tilde{y})) = E(z'u(x, z)).\tag{2}$$

Proof. See Appendix A. ■

Condition (i) defines the set (denoted B) of all observationally equivalent values of parameter β compatible with structural assumptions and the data. Condition (ii) of Theorem

⁵For almost all (v, x, z) in its support, which justifies that we divide by $p_k(x, z)$. Division by zero is a null-probability event.

2 characterizes B , showing that $\beta \in B$ if and only if it satisfies equation (2). As discussed in a Remark in Appendix A, the proof of Theorem 2 also leads to a characterization of the set of observationally equivalent distribution functions $F_\varepsilon(\cdot | x, z)$. Green et al. (1998) proves a special case of this Theorem when neither regressors x nor instruments z are present. It allows them to provide bounds for the average willingness to pay in a contingent valuation experiment.

Before moving on to a more detailed discussion of the characteristics of set B , it is possible to provide a clarifying sketch of its proof by analyzing the case, $K = 2$. Consider (β, F_ε) satisfying Assumption L and its associated reduced form $G_k(x, z)$ for $k = 1, 2$. By Lemma 1 and $(NP2)$, we have $G_1(x, z) = 0$ and $G_2(x, z) = 1$ which makes this case trivial. Restriction $(L.2)$ implies:

$$v_1 \leq -(x\beta + \varepsilon) < v_2$$

When $K = 2$, \tilde{y} is equal to $-(v_2 + v_1)/2$ whatever v and the previous condition can be rewritten:

$$-(v_2 - v_1)/2 \leq -(x\beta + \varepsilon) + \tilde{y} < (v_2 - v_1)/2 = \Delta(x, z)$$

Hence, if we define $u(x, z) = -E(\tilde{y} - x\beta - \varepsilon | x, z)$, it belongs to $] -\Delta(x, z), \Delta(x, z)[$ and satisfies (2), as stated by Theorem 2.

Reciprocally, assume that there exists $u(x, z)$ in $] - (v_2 - v_1)/2, (v_2 - v_1)/2[$ which satisfies condition (2). Consider a random variable λ taking values in $]0, 1[$ and such that:

$$E(\lambda | x, z) = \frac{1}{2} + \frac{u(x, z)}{v_2 - v_1}$$

Then consider the random variable, $\varepsilon = -x\beta - \lambda v_1 - (1 - \lambda)v_2$. By construction, it satisfies $v_1 < -(x\beta + \varepsilon) \leq v_2$. Hence, the model (β, F_ε) satisfies $(L.1-L.2)$ and generates $G_1(x, z) = 0$ and $G_2(x, z) = 1$ through (LV) . The only remaining condition to check is $(L.3)$, namely ε is uncorrelated with z . It is shown using condition (2) and the definition of λ .

Figure 1 provides an illustration of the results stated in Theorem 2. Given some (x, z) , the nodes represent the conditional probability distribution $G(v, x, z)$ as a function of the special regressor, v . In this example, v satisfies $(NP.2)$, namely the conditional probability is equal to 0 at the lower bound ($v = -1$) and equal to 1 at the upper bound ($v = 1$). The other observed values are at $v = -.5, 0, 0.5$. By construction, if (β, F_ε) generates G

through (LV) , it satisfies $1 - F_\varepsilon(-x\beta - v \mid x, z) = G(v, x, z)$. Hence, the only compatible distribution functions of the shock ε are such that $1 - F_\varepsilon(-x\beta - v \mid x, z)$ is passing through the nodes at $v = -.5, 0, 0.5$. The only other restrictions are that these distribution functions are non-decreasing within the rectangles between the nodes. An example is reported in the graph but it is only one among many other possibilities. The total surface of the rectangles is given by function $2\Delta(x, z)$ and it measures the degree of our ignorance on the distribution of ε .

The following section builds on Theorem 2 to provide a more detailed description of B , the set of observationally equivalent parameters.

3.3 Bounds on Structural Parameters and Overidentification

This section provides a more detailed description of B , the set of observationally equivalent parameters. We focus on the case where the number of instruments z is equal to the number of variables x (the exogenous case $z = x$ being the leading example). At the end of the section, we will briefly discuss how the results could be extended to the case where the number of instruments z is larger than the number of explanatory variables, x .

3.3.1 Characterizing the Identified Set

When the number of instruments is equal to the number of variables, the assumption that $E(z'x)$ is full rank (*R.ii*) implies that equation (2) has one and only one solution in β for any function $u(x, z)$. Because equation (2) is linear in β , the set B is convex. Also it is non-empty, since it necessarily contains the focal (say) value β^* associated with the moment condition, $E(z'(x\beta^* - \tilde{y})) = 0$ when $u(x, z) = 0$.

The set B can be described as a neighborhood of β^* which size depends on the distances $(v_k - v_{k-1})$ between the different elements of the support of v . Specifically, β^* can be interpreted as the specific value that β would take if these distances were negligible. First, Theorem 2 makes possible to obtain very simple upper bounds for the potential bias that affects the result of the IV regression of \tilde{y} on x . Denoting the half-length of the largest interval as

$$\Delta_M = \max_{k \in \{2, \dots, K\}} (v_k - v_{k-1}) / 2,$$

we have:

Corollary 3 *The identification set B is non empty and convex. It contains the focal value β^* defined as:*

$$\beta^* = E(z'x)^{-1}E(z'\tilde{y})$$

and any $\beta \in B$ satisfies,

$$(\beta - \beta^*)'W(\beta - \beta^*) \leq E[(\Delta(x, z))^2] \leq \Delta_M^2,$$

where $W = E(x'z)(E(z'z))^{-1}E(z'x)$.

Proof. See Appendix A. ■

Corollary 3 shows that B lies within an ellipsoid whose size is bounded by Δ_M in the metric W . A sharp but less straightforward result will be given in Section 3.3.3. Notice that in the specific case where the different v_k are equidistant (i.e., $\forall k = 3, \dots, K, v_k - v_{k-1} = v_2 - v_1$), the half-length between two successive points, $\Delta_M = \frac{v_2 - v_1}{2}$, provides an upper bound for the size of the ellipsoid.

Returning to the general case, the maximum-length index, Δ_M , can be taken as a measure of distance to continuity of the distribution function of v (or of its support Ω_v). For a latent model $(\beta, F_\varepsilon(\cdot | x, z))$, corollary 3 proves that, for a sequence of support Ω_v indexed by Δ_M , we have:

$$\lim_{\Delta_M \rightarrow 0} B = \{\beta^*\},$$

and point identification is restored.

3.3.2 Interval Identification in the Coordinate Dimensions

This section characterizes the identification region of the different individual parameters. Let

$$B_p = \{\beta_p \in \mathbb{R} \mid \exists(\beta_1, \dots, \beta_{p-1}) \in \mathbb{R}^{p-1}, (\beta_1, \dots, \beta_{p-1}, \beta_p) \in B\}$$

represents the identification region of the last coefficient (say). All scalars belonging to this interval, are observationally equivalent to the p th component of the true parameter.

Corollary 4 *B_p is an interval centered at β_p^* , the p -th component of β^* . Specifically, we have,*

$$B_p = \left] \beta_p^* - \frac{E(|\tilde{x}_p| \Delta(x, z))}{E(\tilde{x}_p^2)}; \beta_p^* + \frac{E(|\tilde{x}_p| \Delta(x, z))}{E(\tilde{x}_p^2)} \right]$$

where \tilde{x}_p is the remainder of the IV-projection of x_p onto the other components of x using instruments z (as formally defined in the proof).

Proof. See Appendix A. ■

Generally speaking, the estimation of B_p requires the estimation of $E(|\tilde{x}_p| \Delta(x, z))$. Given this fact, it is worth emphasizing that $\Delta(x, z)$ can be rewritten $E(\tilde{y}_\Delta | x, z)$ where $\tilde{y}_\Delta = \frac{\mu_k \cdot y}{p_k(x, z)} + \frac{v_K - v_{K-1}}{2}$, with $\mu_k = \frac{(v_k - v_{k-1} - (v_{k+1} - v_k))}{2}$ for $k = 2, \dots, K - 1$ and $\mu_1 = \mu_K = 0$ (as shown at the end of the proof of corollary 5). Hence, $E(|\tilde{x}_p| \Delta(x, z))$ can be rewritten $E(|\tilde{x}_p| \tilde{y}_\Delta)$ which means that the estimation of the upper and lower bounds of B_p only requires [1] the construction of the transform \tilde{y}_Δ , [2] an estimation of the residual \tilde{x}_p and [3] the linear regression of \tilde{y}_Δ on $|\tilde{x}_p|$.

3.3.3 Interval Identification of Linear Combinations of the Parameters

This section characterizes the identification region of any linear combination of the parameters.

Assume for simplicity that x is exogenous and consider Q an orthogonal matrix of dimension p . It satisfies $QQ' = I$ and can be written $Q = (Q_{-p}, q_p)$ where q_p is a vector of dimension p and Q_{-p} a matrix of dimension $[p, (p - 1)]$. Also, we can always write:

$$x\beta = xQQ'\beta = s\beta_Q$$

where $s = xQ$ and $\beta_Q = Q'\beta$. The p -th component of β_Q is, $\beta_Q^{(p)} = q_p'\beta$. It is associated to the p -th explanatory variable, $s_p = xq_p$. Conversely, as Q can be any orthogonal matrix, any linear combination of parameter β can be written as $\beta_Q^{(p)}$ up to a scale parameter which is equal to its Euclidean norm in R^p . Hence, the problem of identifying the vector of parameters β generating (y, v, x, z) through model (LV) is equivalent to the problem of identifying the vector of parameters β_Q generating (y, v, s, z) through the same latent model. Let B_Q be the identification region of β_Q and $B_Q^{(p)}$ the projection of B_Q onto its p -th dimension. Denoting \tilde{s}_p the remainder of the projection of s_p onto $s_{-p} = xQ_{-p}$, we can apply corollary 4 and write,

$$B_Q^{(p)} = [q_p'\beta^* - \frac{E(|\tilde{s}_p| \Delta(x))}{E(\tilde{s}_p^2)}, q_p'\beta^* + \frac{E(|\tilde{s}_p| \Delta(x))}{E(\tilde{s}_p^2)}]$$

For any normalized vector of weights q , this equation provides us with an analytical definition of the set of all the scalars that are observationally equivalent to the true $q'\beta$.

Note that the length of the interval of identification $B_Q^{(p)}$ can be easily expressed as a function of x and Q . Specifically, we have,

$$\begin{aligned}\tilde{s}_p &\equiv s_p - s_{-p} [E(s'_{-p}s_{-p})]^{-1} E(s'_{-p}s_p) \\ &= xq_p - xQ_{-p} [Q'_{-p}E(x'x)Q_{-p}]^{-1} Q'_{-p}E(x'x)q_p, \\ &= x(I - Q_{-p} [Q'_{-p}E(x'x)Q_{-p}]^{-1} Q'_{-p}E(x'x))q_p, \\ &= xM_Q^{(p)}q_p \text{ (say),}\end{aligned}$$

which yields:

$$\frac{E(|\tilde{s}_p | \Delta(x))}{E(\tilde{s}_p^2)} = \frac{E(|xM_Q^{(p)}q_p | \Delta(x))}{q'_pM_Q^{(p)'}E(x'x)M_Q^{(p)}q_p}$$

3.3.4 The Construction of the Identification Region

For any vector q of the unit sphere of \mathbb{R}^p i.e. $\|q\|=1$, denote B_q the set of all scalars that are observationally equivalent to the true $q'\beta$. We know from the previous subsection that B_q is an interval which center is $q'\beta^*$. We are now going to show how to reconstruct the identification region B from all $\{B_q\}_{\|q\|=1}$.

To begin with, let us define $I_q(B)$ the set of parameters β which projections on q are observationally equivalent to the true $q'\beta$,

$$I_q(B) = \{\beta \in \mathbb{R}^p : q'\beta \in B_q\}.$$

For any vector q , $\|q\|=1$, $I_q(B)$ is a band of \mathbb{R}^p which contains B . The intersection of all $I_q(B)$ also contains B . Put differently,

$$B \subset I(B) = \cap_{\|q\|=1} I_q(B).$$

Because B is bounded and convex, we can actually show that $I(B)$ is equal to the identification region B .

Lemma 5 $B = I(B)$.

Proof. See Appendix A. ■

Even if B has no analytical definition equation, it can be constructed very easily by simulation. Randomly draw S vectors q_s , $\|q_s\|=1$, construct the bands $I_{q_s}(B)$ and their intersection $\cap_{s=1,..,S} I_{q_s}(B)$. Then make S go to infinity.

Regarding inference, frameworks of Horowitz and Manski (1998), Imbens and Manski (2004) or Chernozhukov et al. (2004) can be immediately applied to these results. In particular, Chernozhukov et al. (2004) use interval identification as one of their leading examples. Tests and confidence intervals can be readily derived using the objective function that they propose. It is more difficult to extend their results to many dimensions as in our general case above and we left it for future research.

3.3.5 Supplementary Restrictions

A potentially interesting development of this framework is when the number of instruments is larger than the number of variables ($q > p$). In such a case, B is not necessarily non-empty since condition (2) in Theorem 2 may have no solutions at all (i.e., some "overidentification" restrictions may be not true).

Consider z_A , a random vector which dimension is the same as random vector x , defined by:

$$z_A = zA$$

and such that $E(z'_A x)$ is full rank. Define the set, \mathcal{A} , of such matrices A of dimension p, q . The previous analysis can then be repeated for any A in such a set. The identification set $B(A)$ is now indexed by A . Under the maintained assumption (L.3), the true parameter (or parameters) belongs to the intersection of all such sets when matrix A varies:

$$B \subset \bigcap_{A \in \mathcal{A}} B(A)$$

As previously, the set on the RHS is convex because it is the intersection of convex sets. What changes is that it can be empty which refutes the maintained assumption (L.3). This argument would form the basis for optimizing the choice of A or for constructing test procedures of overidentifying restrictions in such a partial identification framework. The question is open whether the usual results hold. Finally, we can always project this set onto its elementary dimensions. The intersection of the projections is the projection of the intersections.

For the sake of simplicity, we shall proceed in the rest of the paper using the assumption that $p = q$ which is worthwhile investigating first.

3.4 Priors on The Range of Variation

Theorem 2 and its corollaries characterize the set of parameters (denoted B) that are observationally equivalent to the true parameter under the assumption that the conditional probability $Pr(y = 1 | v, x, z)$ increases from 0 to 1 when v varies over its support. This condition represents a potentially important limitation in empirical applications. A more careful look at Theorem 2 shows that it is possible to relax this assumption and to characterize the identification set in a substantially more general framework.

Because of (NP.2), one key aspect of Theorem 2 is knowing $Pr(v = v_1 | x, z)$ or $Pr(v = v_K | x, z)$ does not provide any additional information on the identification region. In fact, B can be identified even in the extreme case where $Pr(v = v_1 | x, z) = Pr(v = v_K | x, z) = 0$, when v_1 and v_K are outside the true support of v . Put differently, is not necessary to actually observe $Pr(y = 1 | v, x, z)$ varying from zero to one to identify the set B , it is only necessary to impose this condition as a prior on the data generating process of y at values of v that are not observed in the available data. Some examples are given below.

To be more specific, consider the following reformulation of (L.2),

(L.2bis) There exist two finite real numbers v_0 and v_{K+1} , with $v_0 < v_1$ and $v_{K+1} > v_K$, such that the support of $-x\beta - \varepsilon$ is included in $[v_0, v_{K+1}[$.

Condition (L.2bis) clearly relaxes condition (L.2). Under (L.2bis), $Pr(y = 1 | v, x, z)$ does not necessarily vary from zero to one when v varies over its support $\Omega_v = \{v_1, \dots, v_K\}$, so that $Pr(y = 1 | v, x, z)$ does not necessarily satisfy condition (NP.2) anymore. Condition (L.2bis) imposes (NP.2) as a prior on $Pr(y = 1 | v, x, z)$ for values of v , v_0 and v_{K+1} , that are actually not observed in the data.

It is straightforward to check that B can be identified under (L.2bis) following exactly the same route as under (L.2). The only change is to replace v_1 by v_0 and v_K by v_{K+1} .

Corollary 6 Consider β a vector of parameter and $Pr(y = 1 | v = v_k, x, z)$, a conditional probability distribution (denoted $G_k(x, z)$) satisfying the monotonicity condition (NP.1). The

two following statements are equivalent,

(i) there exists a latent random variable ε such that the latent model $(\beta, F_\varepsilon(\cdot | x, z))$ satisfies conditions (L.1 – L.2bis – L.3) and such that $\{G_k(x, z)\}_{k=1,..,K}$ is its image through the transformation (LV),

(ii) there exists a measurable function $u(x, z)$ from $S_{x,z}$ to \mathbb{R} which takes its values in the interval (a.e. $F_{x,z}$)

$$I(x, z) =] - \Delta(x, z), \Delta(x, z)],$$

where

$$\Delta(x, z) = \frac{1}{2} \sum_{k=1}^{K+1} [(v_k - v_{k-1})(G_k(x, z) - G_{k-1}(x, z))],$$

and such that,

$$E(z'(x\beta - \tilde{y})) = E(z'u(x, z)). \quad (3)$$

This corollary states that identification remains possible even when the support of the special regressor is not large and when the probability of observing $y = 1$ does not vary from zero to one. The cost is that the identification set depends on priors (i.e, v_0 and v_{K+1}) which location might be debatable and the case should be argued in each particular application.

An example of potential application is the analysis of the probability of buying an object (a bottle of water, say) as a function of an experimentally-set price v . Specifically, each individual is faced with a price which is under experimental control and can take only two values v_1 and v_2 . Though we only observe two prices, we can plausibly assume that for a sufficiently small (large) v_0 (v_3) the probability of buying the object is 1 (0) whatever the characteristics of the individuals. Hence, the problem can be redefined with the support of v being $\{v_0, v_1, v_2, v_3\}$ and with the additional assumption that $Pr(y = 1 | v, x, z)$ varies from zero to one when v varies over its support.

Other (non-experimental) examples include the analysis of the probability of entry (or exit) into such basic institutions as the labor market or the school system. Consider for instance the school-leaving probability in a typical developed country where v stands for individuals' age at the end of the year. We can plausibly speculate that the complete variability of this probability between 0 and 1 (Condition NP.2) is satisfied when (say) $v_0 = 15$ years and $v_{K+1} = 30$ years. Using these priors and assuming that the school-leaving

latent propensity may be written $(x\beta + v + \varepsilon)$, we can provide valuable inference on β even if our sample of observations consists in 20 to 25 years old individuals and such that the observed probability of school leaving of the 20 (25) years' old is strictly greater (lower) than 0 (1)⁶. Generally speaking, the identification method developed in this paper can also be applied to the analysis of virtually any binary phenomena which diffusion may be assumed dependent of a secular time-trend.

4 Interval Data

In this section, we deal with the case where v is continuous although it is observed by intervals only. We show that the set of parameters observationally equivalent to the true structural parameter has a similar structure as in the discrete case. It is a convex set and, when there are no overidentifying restrictions ($p = q$), it is not empty. It contains the value corresponding to an IV regression of a transformation of y on x given instruments z . When some information is available on the conditional distribution function of regressor v within-intervals, the identification set shrinks. Its size diminishes as the distribution function of the special regressor within intervals becomes closer to uniformity. When v is conditionally uniformly distributed within intervals, the identification set is a singleton and the parameter of interest β is exactly identified.

4.1 Identification Set: the General Case

The data are now characterized by a random variable (y, v, v^*, x, z) where v^* is the result of censoring v by interval. Only realizations of (y, v^*, x, z) are observed and those of v are not. Variable v^* is discrete and defines the interval in which v lies. More specifically, assumption D is replaced by:

Assumption ID:

(i) (*Interval Data*) The support of v^* conditional on (x, z) is $\{1, \dots, K - 1\}$ almost everywhere $F_{x,z}$. The distribution function of v^* conditional on (x, z) is denoted $p_{v^*}(x, z)$. It is defined almost everywhere $F_{x,z}$.

⁶Under slightly different structural assumptions, this example can also be used in the section dealing with interval data when age is treated as a censored continuous variable.

(ii) (*Continuous Regressor*) The support of v conditional on $(x, z, v^* = k)$ is $[v_k, v_{k+1}[$ (almost everywhere $F_{x,z}$). The overall support is $[v_1, v_K[$. The distribution function of v conditional on x, z, v^* is denoted $F_v(\cdot | v^*, x, z)$ and is assumed to be absolutely continuous. Its density function denoted $f_v(\cdot | v^*, x, z)$ is strictly positive and bounded.

We consider latent models which satisfy the large support condition (L.2) (i.e., the support of $-x\beta - \epsilon$ is included in the support of v), the moment condition (L.3) (i.e., $E(z'\epsilon) = 0$) and the following extension of the partial independence hypothesis,

$$F_\epsilon(\cdot | v, v^*, x, z) = F_\epsilon(\cdot | x, z) \quad (\text{L.1}^*)$$

The conditional probability distributions $Pr(y = 1 | v^*, x, z)$ generated through transformation (LV) by such latent models is non decreasing in v^* using the same argument as in Lemma 1. Distribution $Pr(y = 1 | v^*, x, z)$ does not vary from 0 to 1 however when v^* varies over its support. Using the terminology of the previous section, it necessarily satisfies condition (NP.1), but does not necessarily satisfy condition (NP.2). We will keep the former restriction and drop the latter from the definition of the class of binary reduced forms under consideration.

Interval censorship that we consider does not cover cases where intervals are unbounded on the left and/or on the right. Using Magnac and Maurin (2004), we can indeed prove that parameter β does not belong to a bounded set in these cases. We thus restrict ourselves to this set of assumptions which is, not only in this respect, quite similar to assumptions proposed by Manski and Tamer (2002). Their Interval (I) assumption is equivalent to Assumption (ID.ii), their Monotone assumption (M) is exactly (NP.1) and their Mean Independence assumption (MI) is a consequence of our assumption (L.1*), ours being slightly stronger. We however depart from the quantile restriction and exogeneity assumptions that they use in the binary case since we assume the absence of correlation of shocks with some instruments z . We also depart from their setting since the bounds of the intervals of observation are not random in our case.⁷

We now consider a conditional probability function $Pr(y = 1 | v^*, x, z)$ which is non

⁷We discuss in Magnac and Maurin (2004) why quantile restrictions as in Manski (1985) and partial independence and large support assumptions as in Lewbel (2000) lead to non nested models which attractiveness depends on the context.

decreasing in v^* and we search for a latent model generating this reduced form through transformation (LV). In analogy with the discrete case, we begin with constructing a transformation of the dependent variable. If $\delta(v^*) = v_{v^*+1} - v_{v^*}$ denotes the length of the v^* th interval, the transformation adapted to interval data is :

$$\bar{y} = \frac{\delta(v^*)}{p_{v^*}(x, z)} y - v_K \quad (4)$$

It is slightly different from the transformation (1) in terms of weights $\delta(v^*)$ and in reference to the end-points but the dependence on the random variable $y/p_{v^*}(x, z)$ remains the same.

With these notations, the following theorem analyses the degree of underidentification of the structural parameter β .

Theorem 7 *Consider β a vector of parameter and $Pr(y = 1 | v^*, x, z)$ (denoted $G_{v^*}(x, z)$) a conditional distribution function which is non decreasing in v^* . The two following statements are equivalent,*

(i) *there exist a latent conditional distribution function of v , $F_v(. | x, z, v^*)$, and a latent random variable ε defined by its conditional distribution function $F_\varepsilon(. | x, z)$ such that:*

a. *$(\beta, F_\varepsilon(. | x, z))$ satisfies (L.1*, L.2, L.3)*

b. *$G_{v^*}(x, z)$ is the image of $(\beta, F_\varepsilon(. | x, z))$ through the transformation (LV),*

(ii) *there exists a function $u^*(x, z)$ taking its values in $I^*(x, z) =]\underline{\Delta}^*(x, z), \overline{\Delta}^*(x, z)[$ where (by convention, $G_0(x, z) = 0, G_K(x, z) = 1$),*

$$\begin{aligned} \overline{\Delta}^*(x, z) &= \sum_{k=1, \dots, K-1} (G_{k+1}(x, z) - G_k(x, z))(v_{k+1} - v_k), \\ \underline{\Delta}^*(x, z) &= - \sum_{k=1, \dots, K-1} (G_k(x, z) - G_{k-1}(x, z))(v_{k+1} - v_k), \end{aligned}$$

and such that,

$$E(z'(x\beta - \bar{y}) = E(z'u^*(x, z))). \quad (5)$$

Proof. See Appendix B ■

The identification set has the same general structure in the interval-data case as in the discrete case. It is a non-empty convex set which contains the focal value corresponding to the moment condition $E(z'(x\beta^* - \bar{y})) = 0$.

4.2 Inference Using Additional Information on the Distribution Function of the Special Regressor

We now study how additional information helps to shrink the identification set. There are many instances where there exists additional information on the conditional distribution function of v within intervals. It may correspond to the case where v is observed at the initial stage of a survey or a census, but then dropped from the files that are made available to researchers for confidentiality reasons. Only interval-data information and information (estimates for instance) about the conditional distribution function of v remains available. This framework may also correspond to the case where the conditional distribution function of v is available in one database that does not contain information on y while the information on y is available in another database⁸ which contains only interval information on v .

To analyse these situations, we complete the statistical model by assuming that we have full information on the conditional distribution of v :

(NP.3) : The conditional distribution function of v is known and denoted $\Phi(v | x, z, v^*)$.

The first question is whether this additional information reduces the identification set. The second question is whether there exists an optimal way of censoring v and choosing the intervals for defining v^* . Knowing how identification is related to the conditional distribution $\Phi(v | x, z, v^*)$ may provide interesting guidelines to control censorship.

The first unsurprising result is that additional knowledge on $\Phi(v | x, z, v^*)$ actually helps to shrink the identification set. The second - more surprising - result is that point-identification is restored provided that the conditional distribution function of the censored variable v is piece-wise uniform.

To state these two results, we are going to use indexes measuring the distance of a distribution function $\Phi(v | v^* = k, x, z)$ to uniformity. Specifically, we denote $U(v | v^* = k) = \frac{v - v_k}{v_{k+1} - v_k}$ the uniform c.d.f and we consider the two following indexes,

⁸Angrist and Krueger (1992) or Arellano and Meghir (1992) among others developed two-sample IV techniques for such data design in the linear case.

$$\begin{aligned}\xi_k^U(x, z) &= \sup_{v \in]v_k, v_{k+1}[} \left[\frac{\Phi - U}{\Phi} \right] \\ \xi_k^L(x, z) &= \inf_{v \in]v_k, v_{k+1}[} \left[\frac{\Phi - U}{1 - \Phi} \right]\end{aligned}$$

where the arguments of Φ and U are made implicit for expositional simplicity.

Given that Φ is absolutely continuous and its density is positive everywhere (*ID(ii)*), $\frac{\Phi-U}{\Phi}$ and $\frac{\Phi-U}{1-\Phi}$ are well defined on $]v_k, v_{k+1}[$ and satisfy $\frac{\Phi-U}{\Phi} < 1$ and $\frac{\Phi-U}{1-\Phi} > -1$. Furthermore, given that $\frac{\Phi-U}{\Phi}(\frac{\Phi-U}{1-\Phi})$ is continuous and equal to zero at v_{k+1} (at v_k), the supremum of this function in the neighborhood of $v_{k+1}(v_k)$ is clearly non negative (non positive). Hence, we have $\xi_k^L(x, z) \in]-1, 0]$ and $\xi_k^U(x, z) \in [0, 1[$, the two indices being equal to zero when Φ is equal to U . Using these notations, we have the following theorem:

Theorem 8 *Consider β a vector of parameters, $Pr(y = 1 \mid v^*, x, z)$ (denoted $G_{v^*}(x, z)$) a conditional distribution function which is non decreasing in v^* and $\Phi(v \mid v^*, x, z)$ a conditional distribution function. The two following statements are equivalent,*

(i) *there exists a latent random variable ε defined by its conditional distribution function $F_\varepsilon(\cdot \mid x, z)$ such that:*

- a. *$(\beta, F_\varepsilon(\cdot \mid x, z))$ satisfies (L.1*, L.2, L.3)*
- b. *$G_{v^*}(x, z)$ is the image of $(\beta, F_\varepsilon(\cdot \mid x, z))$ through the transformation (LV),*

(ii) *there exists a function $u^*(x, z)$ taking its values in $[\underline{\Delta}_\Phi^*(x, z), \overline{\Delta}_\Phi^*(x, z)]$ where:*

$$\begin{aligned}\underline{\Delta}_\Phi^*(x, z) &= \sum_{k=1, \dots, K-1} (v_{k+1} - v_k) \xi_k^L(x, z) (G_k(x, z) - G_{k-1}(x, z)) \\ \overline{\Delta}_\Phi^*(x, z) &= \sum_{k=1, \dots, K-1} (v_{k+1} - v_k) \xi_k^U(x, z) (G_{k+1}(x, z) - G_k(x, z))\end{aligned}$$

and such that,

$$E(z'(x\beta - \bar{y}) = E(z'u^*(x, z))).$$

Proof. See Appendix B ■

Given that $\xi_k^L(x, z) \in]-1, 0]$ and $\xi_k^U(x, z) \in [0, 1[$, the identification set characterized by Theorem 8 is clearly smaller than the identification set characterized by Theorem 7 when no information is available on v . Also, Theorem 8 makes clear that the size of identification

set diminishes with respect to the distance between the conditional distribution of v and the uniform distribution, as measured by $\xi_k^L(x, z)$ and $\xi_k^U(x, z)$. When this distance is abolished and v is piece-wise uniform, the identification set boils down to a singleton.

Corollary 9 *The identification set is a singleton if and only if the conditional distribution, $\Phi(v | x, z, v^*)$, for all $v^* = k$, and a.e. $F_{x,z}$, is uniform, i.e.:*

$$\Phi(v | v^* = k, x, z) = \frac{v - v_k}{v_{k+1} - v_k}$$

Proof. See Appendix B ■

Corollary 9 corresponds to the “best” case and an interpretation of this result is offered at the end of the proof in Appendix B. It is very different from sufficient conditions for identification given in Manski and Tamer (2002, Corollary page 524). Their conditions (c) and (d) for point identification imply that the probability that the interval of observation of v (denoted v_0, v_1 in their notations) is as small as we want, is positive. In our case, this length is fixed. As our condition is necessary and sufficient, a complete comparison with what can be obtained in the setting of Manski and Tamer (2002) is out of the scope of this paper.

Assuming that the distribution of v is not piece-wise uniform, the question remains whether it is possible to rank the potential distributions of v according to the corresponding degree of underidentification of β . The answer is positive. Specifically, the closer to uniformity the conditional distribution of v is, the smaller the identification set is.

To state this result, we first need to rank distributions according to the magnitude of their deviations from the uniform distribution.

Definition 10 $\Phi_2(v | x, z, v^*)$ is closer to uniformity than $\Phi_1(v | x, z, v^*)$, when a.e. $F_{x,z}$ and for any $k \in \{1, \dots, K - 1\}$:

$$\begin{aligned} \xi_{k,1}^L(x, z) &\leq \xi_{k,2}^L(x, z) \\ \xi_{k,1}^U(x, z) &\geq \xi_{k,2}^U(x, z). \end{aligned}$$

The corresponding preorder is denoted $\Phi_1 \succeq \Phi_2$.

Using this definition:

Corollary 11 *Let $\Phi(v \mid v^* = k, x, z)$ any conditional distribution. Let B the associated region of identification for β . Then:*

$$\Phi_1 \succeq \Phi_2 \implies B_{\Phi_2} \subseteq B_{\Phi_1}$$

Proof. Straightforward using Theorem 8. ■

Assuming that we have some control on the construction on v^* (i.e., on the information on v that are made available to researchers), this result shows that it has simply to be constructed in a way that minimizes the distance between the uniform distribution and the distribution of v conditional on v^* (and other regressors). Such a choice minimizes the length of the identification interval. Consider for instance date of birth. The frequency of this variable plausibly varies from one season to another, or even from one month to another, especially in countries where there exist strong seasonal variations in economic activity. At the same time, it is likely that the frequency of this variable does not vary significantly within months, meaning it is likely that it is uniformly distributed within months in most countries. In such a case, our results show that we only have to made available the month of birth of respondents (and not necessarily their exact date-of-birth) to achieve exact identification of structural parameters of binary models which are monotone with respect to date-of-birth.

4.3 Projections of the Identification Set

Results concerning projections of the identification set in the discrete case can be easily extended to the case of interval data. As in the discrete case and for simplicity, we restrict our analysis to the leading case when the dimension of z and x are the same. The identified set B can be projected onto its elementary dimensions using the same usual rules as in Corollary 4.

Let:

$$B_p = \{\beta_p \in \mathbb{R} \mid \exists(\beta_1, \dots, \beta_{p-1}) \in \mathbb{R}^{p-1}, (\beta_1, \dots, \beta_{p-1}, \beta_p) \in B\}$$

be the projected set corresponding to the last (say) coefficient. All scalars belonging to this interval, are observationally equivalent to the p th component of the true parameter. We denote β^* the solution of equation (5) when function $u^*(x, z) = 0$ (as $E(z'x)$ is a square

invertible matrix):

$$\beta^* = E(z'x)^{-1}E(z'\bar{y})$$

To begin with, we consider the case where no information is available on the distribution of v and state the corollary to Theorem 7.

Corollary 12 B_p is an interval which center is β_p^* , where β_p^* represents the p -th component of β^* . Specifically, we have,

$$B_p =]\beta_p^* + \varsigma_{L,p}; \beta_p^* + \varsigma_{U,p}]$$

where :

$$\varsigma_{L,p} = [E(\tilde{x}_p^2)]^{-1} E(\tilde{x}_p(\mathbf{1}\{\tilde{x}_p > 0\}\underline{\Delta}^*(x, z) + \mathbf{1}\{\tilde{x}_p \leq 0\}\bar{\Delta}^*(x, z)))$$

$$\varsigma_{U,p} = [E(\tilde{x}_p^2)]^{-1} E(\tilde{x}_p(\mathbf{1}\{\tilde{x}_p \leq 0\}\underline{\Delta}^*(x, z) + \mathbf{1}\{\tilde{x}_p > 0\}\bar{\Delta}^*(x, z)))$$

with \tilde{x}_p is the residual of the projection of x_p onto the other components of x using instruments z .

Proof. See Appendix B. ■

Corollary 12 has exactly the same structure as the corresponding corollary to Theorem 8, with $\underline{\Delta}^*$ and $\bar{\Delta}^*$ replacing $\underline{\Delta}_{\Phi}^*$ and $\bar{\Delta}_{\Phi}^*$.

Generally speaking, the estimation of B_p requires the estimation of $\varsigma_{L,p}$ and $\varsigma_{U,p}$. At the end of the proof of Corollary 12, we show that these scalars can be estimated through simple regressions. Specifically, let us denote $\bar{y}_L = \frac{\theta_{L,v^*} \cdot y}{p_k(x, z)} + v_K - v_{K-1}$, where $\theta_{L,k} = \frac{(v_{k+2} - v_{k+1} - (v_{k+1} - v_k))}{2}$ for $k = 2, \dots, K-1$ and where $v_{K+1} = v_K$ by convention. Similarly, define $\bar{y}_U = \frac{\theta_{U,v^*} \cdot y}{p_k(x, z)} + v_K - v_{K-1}$, where $\theta_{U,k} = \frac{(v_k - v_{k-1} - (v_{k+1} - v_k))}{2}$ for $k = 2, \dots, K-1$ and where $v_0 = v_1$.

Using these notations, $\varsigma_{L,p}$ is the regression coefficient of $(\mathbf{1}\{\tilde{x}_p > 0\}\bar{y}_{L,p} + \mathbf{1}\{\tilde{x}_p \leq 0\}\bar{y}_{U,p})$ on \tilde{x}_p and $\varsigma_{U,p}$ is the regression coefficient of $(\mathbf{1}\{\tilde{x}_p \leq 0\}\bar{y}_{L,p} + \mathbf{1}\{\tilde{x}_p > 0\}\bar{y}_{U,p})$ on \tilde{x}_p . Interestingly, when all intervals have the same length, \bar{y}_L and \bar{y}_U are equal and constant and the length of the one-dimensional identification region is then proportional to this constant.

5 Estimation and Monte Carlo Experiments

Deriving estimators for the upper and lower bounds of intervals of interest is straightforward since these bounds can be expressed as moments. When the number of observations become large, their properties conform with theoretical properties that have just been derived. It remains to be seen how these estimators of the bounds behave in small and medium-sized samples (i.e., 100 to 1000 observations). This is why we present Monte Carlo experiments in this section. The simulated model is $y = \mathbf{1}\{1 + v + x_2 + \varepsilon > 0\}$. For the sake of clarity, the set-up is chosen to be as close as possible to the set-up originally used by Lewbel (2000). We adapt this original setting to the case where regressor v is interval-valued.⁹

Specifically, the construction of the special regressor v , the covariate x_2 , the instrument z and the random shock ε proceeds in two steps. To begin with, consider four random variables such as: e_1 is uniform on $[0, 1]$, e_2 and e_3 are zero mean unit variance normal variates and e_4 is a mixture of a normal variate $N(-.3, .91)$ using a weight of .75 and a normal variate $N(.9, .19)$ using a weight of .25. Using these notations, we define:

$$\begin{aligned}\eta &= 2e_2 + \alpha e_4, & x_2 &= e_1 + e_4 \\ \varepsilon &= \rho(e_1 - .5) + e_3, & z &= e_4.\end{aligned}$$

where α is a parameter that makes the random shock a non-normal variate and ρ is a parameter that renders x_2 endogenous. The case where $\alpha = \rho = 0$ (resp. $\alpha = \rho = 1$) roughly corresponds to what Lewbel calls the simple (resp. messy) design.

Regressor v is finally defined by truncating η to the 95% symmetric interval around 0, denoted $[v_1, v_K]$. To comply with assumption *L.2*, we then truncate $x_2 + \varepsilon$ so that $1 + x_2 + \varepsilon + v_1 > 0$ and $1 + x_2 + \varepsilon + v_K < 0$. We construct the censored $K - 1$ intervals in the obvious way:

$$v^* = k \text{ if } v \in [v_k, v_{k+1}[$$

5.1 Statistical Summaries

Tables 1 to 4 report various Monte Carlo experiments where the sample size varies using 100, 200, 500 or 1000 observations. In all experiments, the number of Monte Carlo replications

⁹Estimates in the case where v is discrete exhibit very similar properties. Those results are available upon request to the authors.

is equal to 1000. Additional replications do not affect any estimates (resp. standard errors) by more than a 1% margin of error (resp. 3%). We report results in two panels. In the top panel, we report estimates of the lower and upper bounds of both coefficients (intercept and variable) by recentering them at zero instead of their true values which are equal to one. In the bottom panel, we compute the average of the estimates of the lower and upper bounds, $E(\hat{\theta}_b + \hat{\theta}_u)/2$, the adjusted length of the interval, $E(\hat{\theta}_u - \hat{\theta}_b)/2\sqrt{3}$, and the average sampling error defined as:

$$(\hat{\sigma}_u^2 + \hat{\sigma}_b^2 + \hat{\sigma}_u \hat{\sigma}_u)/3$$

where $\hat{\sigma}_u$ and $\hat{\sigma}_b$ are estimated standard errors of the estimated lower and upper bounds. These three statistics actually provide an interesting decomposition of the mean square error uniformly integrated over the interval $[\hat{\theta}_b, \hat{\theta}_u]$:

$$\begin{aligned} MSEI &= E \int_{\hat{\theta}_b}^{\hat{\theta}_u} (\theta - \theta_0)^2 \frac{d\theta}{\hat{\theta}_u - \hat{\theta}_b} \\ &= \frac{1}{3} E \left[\frac{(\hat{\theta}_u - \theta_0)^3 - (\hat{\theta}_b - \theta_0)^3}{\hat{\theta}_u - \hat{\theta}_b} \right] \\ &= \frac{1}{3} E \left[(\hat{\theta}_u - \theta_0)^2 + (\hat{\theta}_b - \theta_0)^2 + (\hat{\theta}_b - \theta_0)(\hat{\theta}_u - \theta_0) \right] \end{aligned}$$

Let $\bar{\theta}_i = E(\hat{\theta}_i)$, $i = u, b$, the expected values of the estimates, and $\bar{\theta}_m = (\bar{\theta}_u + \bar{\theta}_b)/2$ the average center of the interval. We then have:

$$\begin{aligned} MSEI &= (\bar{\theta}_m - \theta_0)^2 + \frac{1}{3} E \left[(\hat{\theta}_u - \bar{\theta}_m)^2 + (\hat{\theta}_b - \bar{\theta}_m)^2 + (\hat{\theta}_b - \bar{\theta}_m)(\hat{\theta}_u - \bar{\theta}_m) \right] \\ &= (\bar{\theta}_m - \theta_0)^2 + \frac{1}{3} \left((\bar{\theta}_u - \bar{\theta}_b)/2 \right)^2 \\ &\quad \frac{1}{3} E \left[(\hat{\theta}_u - \bar{\theta}_u)^2 + (\hat{\theta}_b - \bar{\theta}_b)^2 + (\hat{\theta}_b - \bar{\theta}_b)(\hat{\theta}_u - \bar{\theta}_u) \right] \end{aligned}$$

The first term is the square of a “decentering” term which can be interpreted as a bias term. The second term is the square of the “adjusted” length, which can be interpreted as the “uncertainty” due to partial identification instead of point identification. The third term is an average of standard errors and can then be interpreted as sample variability. These three terms are reported in the bottom panel for both coefficients as well as root mean square error, $MSEI^{1/2}$.

5.2 Results

In Tables 1 to 4, we make some parameters vary: The bandwidth used to compute the denominator of transformation (4) in Table 1, the degree of non normality in Table 2, the degree of endogeneity in Table 3 and the number of points in the support of the special regressor in Table 4. In all cases, the true value of the parameter belongs to the interval built up around the estimates of the lower and upper bounds. Horowitz and Manski (2000) and Imbens and Manski (2004) for an alternative, rigorously define confidence intervals when identification is partial. We here report confidence intervals for bounds only. In cases where the number of points is fixed (Tables 1 to 3), the stability of the estimated length of the interval across experiments is a noticeable result. It almost never vary by more than a relative factor of 10%.

In Table 1, we experimented with different bandwidths. As said, interval lengths are stable, though intervals can be severely decentered for the intercept term. Increasing the sample size or the bandwidth recenters the interval around the true value. Increasing the bandwidth decenters interval estimates for the coefficient of the variable towards the negative numbers though at a much lesser degree. Finally, the mean square error ($MSEI$) for the intercept decreases with the bandwidth while it has frequently a U-shape form for the coefficient of the variable. We have tried to look for a data-driven choice of the bandwidth by minimizing this quantity but it was inconclusive. A larger bandwidth seems to be always preferred. Some further research is clearly needed on this issue.

In Table 2, we experimented with different degrees of non-normality, by making parameter α vary. If this parameter increases, interval length is very weakly affected. Decentering can be quite severe above all for the coefficient of the variable when the degree of non-normality is large. It is also true at a lesser degree for the intercept.

In Table 3, we experimented with different degrees of correlation between covariates and errors and therefore the amount of endogeneity. It is the only case where interval length sometimes differs across experiments with no clear pattern of increase or decrease. There is also some large decentering of the intervals that mildly decreases with the sample size. Standard errors are affected by the degree of endogeneity only when the sample size is less than 200.

In Table 4, we experimented with varying the number of points of the discrete support. Theory predicts that interval length should decrease with the number of points of support. In our experiments, it is always true and this decrease is not much affected by sample sizes. We obtain that result by estimating the conditional probability function of v using nearest neighbors (w.r.t. v) and using kernels for the other covariates. A preliminary less careful estimation of this probability function led to humps and bumps in the estimates. There can be some strong decentering problems though and there is evidence of a trade-off between the length of the interval and the average standard errors. The latter tend to increase when the number of points in the support increases. No doubt that it is partly due to the way we built up the probability estimates. The adaptation of kernel methods proposed by Racine and Li (2004) could be an alternative to deal with this problem.

6 Conclusion

In this paper, we explored partial identification of coefficients of binary variable models when the very exogenous regressor is discrete or interval-valued. We derived bounds for the coefficients and show that they can be written as moments of the data generating process. We also show that in the case of interval data, additional information can shrink the identification set. When the unknown variable is distributed uniformly within intervals, these sets are reduced to one point.

Some additional points seem to be worthwhile considering. First, we do not provide proofs of consistency and asymptotic properties of the estimates of the bounds of the intervals because they would add little to the ones Lewbel (2000) presents. The asymptotic variance-covariance matrix of the bounds can also be derived along similar lines. Moreover, adapting the proofs of Magnac and Maurin (2004) these estimates are efficient in a semi-parametric sense under some conditions.

Generally speaking, the identification results obtained in this paper when data are not continuous may be used to enhance identification power when the data are actually continuous. Specifically, if the support of the continuous very exogenous regressor is not large enough, one could use additional measurements or priors at discrete points at the left and right of the actual support in order to achieve partial or point identification. Such additional

information generate a case with mixed discrete and continuous support. It can be analyzed by using simultaneously the proofs used in the discrete, interval or continuous settings. An interesting situation corresponds to a binary variable which probability of occurrence is known to be monotone in some regressor v and varies between 0 and 1 in a known interval. School-leaving (as a function of age) is such an example. In such a case, the coefficients of the binary latent model are partially identified regardless of whether the scheme of observation of the very exogenous regressor is complete, discrete, by interval or continuous. Two extreme cases lead to exact identification, i.e. complete and continuous observation in the interval on the one hand, and, on the other hand, complete & interval-data observation when the distribution of the very exogenous regressor is uniform within intervals. Other cases are nevertheless still informative.

REFERENCES

- Angrist, J.D., and A. Krueger**, 1992, "The Effect of Age at School Entry on Educational Attainment : An Application of Instrumental Variables with Moments from Two Samples", *Journal of the American Statistical Association*, 57:11-25.
- Arellano, M. and C., Meghir**, 1992, "Female Labor Supply and On-the-Job Search: An Empirical Model Estimated Using Complementary Datasets", *Review of Economic Studies*, 59:537-560.
- Bierens, H.J., and J., Hartog**, 1988, "Nonlinear Regression with Discrete Explanatory Variables with an Application to the Earnings Function", *Journal of Econometrics*, 38:269-299.
- Chernozhukov, V., H. Hong, E. Tamer**, 2004, "Inference on Parameter Sets in Econometric Models", unpublished manuscript.
- Chesher, A.**, 2003, "Non Parametric Identification under Discrete Variation", *Econometrica*, 73:1525-1550.
- Green, D., K.E. Jacowitz, D. Kahneman, D. McFadden**, 1998, "Referendum Contingent Valuation, Anchoring and Willingness to Pay for Public Goods", *Resource and Energy Economics*, 20:85-116.
- Heckman J.J., and E. Vytlacil**, 2005, "Structural Equations, Treatment Effects and Econometric Policy Evaluation", *Econometrica*, 73:669-738.
- Honoré, B., and A., Lewbel**, 2002, "Semiparametric Binary Choice Panel Data Models without Strict Exogeneity", *Econometrica*, 70:2053-2063.
- Horowitz, J.**, 1998, *Semiparametric methods in Econometrics*, Springer: Berlin.
- Horowitz, J., and C.,F., Manski**, 2000, "Non Parametric Analysis of Randomized Experiments with Missing Covariate and Outcome Data", *Journal of the American Statistical Association*, 95:77-84.
- Imbens, G., and C.F., Manski**, 2004, "Confidence Intervals for Partially Identified Parameters", *Econometrica*, 72:1845-1859.
- Leamer, E.E.**, 1987, "Errors in Variables in Linear Systems", *Econometrica*, 55(4): 893-909.
- Lewbel, A.**, 2000, "Semiparametric Qualitative Response Model Estimation with Unknown Heteroskedasticity or Instrumental Variables", *Journal of Econometrics*, 97:145-77.
- Magnac, T., and E., Maurin**, 2004, "Identification & Information in Monotone Binary Models", forthcoming *Journal of Econometrics*.
- Manski, C.F.**, 1985, "Semiparametric Analysis of Discrete Response: Asymptotic Properties of the Maximum Score Estimator", *Journal of Econometrics*, 27:313-33.
- Manski, C.F.**, 1988, "Identification of Binary Response Models", *Journal of the American Statistical Association*, 83:729-738.
- Manski, C.F.**, 2003, *Partial Identification of Probability Distributions*, Springer-Verlag: Berlin.
- Manski, C.F., and E. Tamer**, 2002, "Inference on Regressions with Interval Data on a Regressor or Outcome", *Econometrica*, 70:519-546.

Powell, J., 1994, "Estimation of Semiparametric Models", in eds. R. Engle and D. McFadden, *Handbook of Econometrics*, 4:2444-2521.

Racine, J. and Q., Li, 2004, "Non Parametric Estimation of Regression Functions with Both Categorical and Continuous Data", *Journal of Econometrics*, 119:99-130.

Takayama, A., 1974, *Mathematical Economics*, Cambridge University Press, Cambridge.

A Proofs in Section 3

A.1 Proof of Theorem 2

Let $\{G_k(x, z)\}_{k=1, \dots, K}$ satisfy (NP.1) and (NP.2). It is an ordered set of functions such that $G_1 = 0$ and $G_K = 1$. Fix β . We first prove that (i) implies (ii).

(Necessity) Assume that there exists a latent random variable ε such that $(\beta, F_\varepsilon(\cdot | x, z))$ satisfies (L.1–L.3) and such that $\{G_k(x, z)\}_{k=1, \dots, K}$ is its image through transformation (LV). By (L.2), the conditional support of ε given (x, z) , is included in $] - (v_K + x\beta), -(v_1 + x\beta)]$ and we can write,

$$\forall k; G_k(x, z) = \int_{-(v_k + x\beta)}^{-(v_1 + x\beta)} f_\varepsilon(\varepsilon | x, z) d\varepsilon = 1 - F_\varepsilon(-(v_k + x\beta) | x, z). \quad (\text{A.1})$$

Put differently, we necessarily have $F_\varepsilon(-(v_k + x\beta) | x, z) = 1 - G_k(x, z)$, for each k in $\{1, \dots, K\}$.

Denote $s_k = (v_k + v_{k-1})/2$ and $\delta_k = \frac{v_{k+1} - v_{k-1}}{2} = s_{k+1} - s_k$ for all $k = 2, \dots, K-1$. Setting $\delta_1 = \delta_K = 0$, the transformed variable \tilde{y} is $(\frac{\delta_k y}{p_k(x, z)} - s_K)$ where $y = \mathbf{1}\{v > -(x\beta + \varepsilon)\}$. Integrate \tilde{y} with respect to v and ε :

$$\begin{aligned} E(\tilde{y} \mid x, z) &= \int_{\Omega(\varepsilon|x,z)} \left[\sum_{k=1}^K \delta_k \mathbf{1}\{v_k > -(x\beta + \varepsilon)\} \right] f(\varepsilon | x, z) d\varepsilon - s_K \\ &= \int_{\Omega(\varepsilon|x,z)} \left[\sum_{k=2}^{K-1} (s_{k+1} - s_k) \mathbf{1}\{v_k > -(x\beta + \varepsilon)\} \right] f(\varepsilon | x, z) d\varepsilon - s_K \end{aligned}$$

As the support of $w = -(x\beta + \varepsilon)$ is included in $[v_1, v_K]$, we can also define an integer function $j(w)$ in $\{1, \dots, K-1\}$, such that $v_{j(w)} \leq w < v_{j(w)+1}$. By construction, $v_k > w \Leftrightarrow k > j(w)$ and $\sum_{k=2}^{K-1} (s_{k+1} - s_k) \mathbf{1}\{v_k > w\} = (s_K - s_{j(-x\beta + \varepsilon)+1})$. Hence, we have :

$$\begin{aligned} E(\tilde{y} \mid x, z) &= \int_{\Omega(\varepsilon|x,z)} (s_K - s_{j(-x\beta + \varepsilon)+1}) f(\varepsilon | x, z) d\varepsilon - s_K = -E[s_{j(-x\beta + \varepsilon)+1} \mid x, z] \\ &= x\beta + E(\varepsilon | x, z) - E[s_{j(-x\beta + \varepsilon)+1} + x\beta + \varepsilon \mid x, z] \\ &= x\beta + E(\varepsilon | x, z) - u(x, z) \end{aligned} \quad (\text{A.2})$$

where (recall that $w = -(x\beta + \varepsilon)$):

$$u(x, z) = E(s_{j(w)+1} - w \mid x, z).$$

Bounds on $u(x, z)$ can be obtained using the definition of $j(w)$. First, given that $v_{j(w)} \leq w < v_{j(w)+1}$, we have :

$$-\frac{v_{j(w)+1} - v_{j(w)}}{2} < \frac{v_{j(w)+1} + v_{j(w)}}{2} - w \leq \frac{v_{j(w)+1} - v_{j(w)}}{2}$$

which yields :

$$-\frac{v_{j(w)+1} - v_{j(w)}}{2} < s_{j(w)+1} - w \leq \frac{v_{j(w)+1} - v_{j(w)}}{2}$$

Hence, we can write using the upper bound,

$$\begin{aligned} E(s_{j(w)+1} - w \mid x, z) &= \sum_{k=2}^K \int_{-(v_k+x\beta)}^{-(v_{k-1}+x\beta)} (s_k + x\beta + \epsilon) f(\epsilon \mid x, z) d\epsilon \\ &\leq \sum_{k=2}^K \int_{-(v_k+x\beta)}^{-(v_{k-1}+x\beta)} \frac{v_k - v_{k-1}}{2} f(\epsilon \mid x, z) d\epsilon \\ &= \sum_{k=2}^K \left[\frac{v_k - v_{k-1}}{2} (G_k(x, z) - G_{k-1}(x, z)) \right] = \Delta(x, z) \end{aligned}$$

where in the last equation, we use equation (A.1). Using the lower bound, the proof is similar and thus:

$$-\Delta(x, z) < u(x, z) \leq \Delta(x, z).$$

Since $G_K(x, z) = 1$ and $G_1(x, z) = 0$, we have $\Delta(x, z) \geq \min_k(\frac{v_k - v_{k-1}}{2})$, meaning that $\Delta(x, z) > 0$ and that $I(x, z)$ is non-empty. It finishes the proof that statement (i) implies statement (ii) since equation (A.2) implies (2).

(*Sufficiency*) Conversely, let us prove that statement (ii) implies statement (i). We assume that there exists $u(x, z)$ in $I(x, z) =] - \Delta(x, z), \Delta(x, z)]$ such that equation (2) holds true and we construct a distribution function $F_\epsilon(\cdot \mid x, z)$ satisfying (L.1 – L.3) such that the image of $(\beta, F_\epsilon(\cdot \mid x, z))$ through (LV) is $\{G_k(x, z)\}_{k=1, \dots, K}$.

First, let λ a random variable which support is $]0, 1]$, which conditional density given (v, x, z) is independent of v (a.e. $F_{x,z}$) and is such that:

$$E(\lambda \mid x, z) = (u(x, z) + \Delta(x, z)) / (2\Delta(x, z)). \quad (\text{A.3})$$

As this expectation lies between 0 and 1, it is always possible to find such a random variable. Second, let κ a discrete random variable which support is $\{2, \dots, K\}$ and which conditional distribution given (v, x, z) is independent of v and is given by:

$$Pr(\kappa = k \mid x, z) = G_k(x, z) - G_{k-1}(x, z). \quad (\text{A.4})$$

For any $k \in \{2, \dots, K\}$, consider $K - 1$ random variables, say $\epsilon(\lambda, k)$ which are constructed from λ by:

$$\epsilon(\lambda, k) = -x\beta - \lambda v_{k-1} - (1 - \lambda)v_k$$

Given that $\lambda > 0$, the support of $\epsilon(\lambda, k)$ is $] - x\beta - v_k, -x\beta - v_{k-1}]$. Finally, consider the random variable:

$$\varepsilon = \epsilon(\lambda, \kappa) \quad (\text{A.5})$$

which support is $] - x\beta - v_K, -x\beta - v_1]$ and which is independent of v (because both λ and κ are). It therefore satisfies (L.1) and (L.2). Furthermore, because of (A.4), the image of

$(\beta, F_\varepsilon(\cdot | x, z))$ through (LV) is $\{G_k(x, z)\}_{k=1,..,K}$ because these functions satisfy equation (A.1). The last condition to prove is (L.3). Consider, for almost any (x, z) ,

$$\begin{aligned}
& \int_{\Omega(\varepsilon|x,z)} (s_{j(-x\beta-\varepsilon)+1} + x\beta + \varepsilon)f(\varepsilon | x, z)d\varepsilon = \\
& \sum_{k=2}^K \left(\int_{-x\beta-v_k}^{-x\beta-v_{k-1}} \left(\frac{v_k + v_{k-1}}{2} + x\beta + \varepsilon \right) f(\varepsilon | x, z, \kappa = k) d\varepsilon \right) \\
& = \sum_{k=2}^K E \left(\frac{v_k + v_{k-1}}{2} - \lambda v_{k-1} - (1 - \lambda)v_k | x, z \right) (G(v_k, x, z) - G(v_{k-1}, x, z)) \\
& = \sum_{k=2}^K E(\lambda - 1/2 | x, z) \cdot (v_k - v_{k-1}) \cdot (G_k(x, z) - G_{k-1}(x, z)) \\
& = (u(x, z)/(2\Delta(x, z)))(2\Delta(x, z)) = u(x, z).
\end{aligned}$$

where the third line is the consequence of the definition of ε and the last line is using equation (A.3). Therefore, equation (A.2) holds and:

$$E(z'\tilde{y}) = E(z'x)\beta + E(z'\varepsilon) - E(z'u(x, z)).$$

Equation (2) implies $E(z'\varepsilon) = 0$, that is (L.3), which finishes the proof of Theorem 2. ■

Remark: It is worth emphasizing that this proof also provides a characterization of the domain of observationally equivalent distribution functions F_ε , i.e. the set of random variables ε such that there exists β with (β, F_ε) satisfying conditions (L.1 – L.3) and generating $\{G_k(x, z)\}_{k=1,..,K}$. We have:

The two following statements are equivalent,

- (i) *there exists a vector of parameter β such that the latent model $(\beta, F_\varepsilon(\cdot | x, z))$ verifies conditions L and such that $\{G_k(x, z)\}_{k=1,..,K}$ is its image through the transformation (LV),*
- (ii) *there exist two independent random variables (λ, κ) , conditional on (x, z) , such that the support of λ is $]0, 1]$, the support of κ is $\{2, .., K\}$, equation (A.4) holds and such that:*

$$\varepsilon = -x\beta - \lambda v_{\kappa-1} - (1 - \lambda)v_\kappa$$

where β verifies:

$$E(z'(x\beta - \tilde{y})) = E(z'\Delta(x, z))(2\lambda - 1))$$

A.2 Proof of Corollary 3

First, B contains β^* because $u(x, z) = 0$ takes its values in the admissible set, $I(x, z)$. Second, B is convex because $I(x, z)$ is convex and equation (2) is linear. Furthermore, assume that $(\beta, F_\varepsilon(\cdot | x, z))$ satisfies condition L and generates $G(v, x, z)$ through the transformation (LV). Using Theorem 2, there exists $u(x, z) \in I(x, z)$ such that,

$$E(z'x)(\beta - \beta^*) = E(z'u(x, z))$$

and thus using the definition of W :

$$(\beta - \beta^*)'W(\beta - \beta^*) = E(u'(x, z)z)E(z'z)^{-1}E(z'u(x, z)).$$

Using the generalized Cauchy-Schwarz inequality, we have,

$$E(u'(x, z)z)E(z'z)^{-1}E(z'u(x, z)) \leq E(u(x, z)^2).$$

and by Theorem 2, $E(u(x, z)^2) \leq E(\Delta(x, z)^2)$. By definition, $E(\Delta(x, z)^2) \leq \Delta_M^2$ which completes the proof.

A.3 Proof of Corollary 4

For the sake of clarity, we start with the exogeneous case where $z = x$. Denote x_p the last variable in x , x_{-p} all the other variables (i.e., $x = (x_{-p}, x_p)$). Consider any $\beta \in B$ and $\beta^* = (E(x'x))^{-1}E(x'y)$. There exists a function $u(x)$ in $]-\Delta(x), \Delta(x)[$ such that $\beta - \beta^* = (E(x'x))^{-1}E(x'u(x))$ which is also the result of the regression of $u(x)$ on x .

Denote the residual of the projection of x_p onto the other components x_{-p} as \tilde{x}_p :

$$\tilde{x}_p = x_p - x_{-p} (E(x'_{-p}x_{-p}))^{-1} E(x'_{-p}x_p)$$

Applying the principle of Frish-Waugh, we have

$$\beta_p - \beta_p^* = (E((\tilde{x}_p)^2))^{-1} E(\tilde{x}_p u(x)),$$

since \tilde{x}_p is a scalar. The maximum (resp. minimum) of $E(\tilde{x}_p u(x))$ when $u(x, z)$ varies in $]-\Delta(x), \Delta(x)[$ is obtained by setting $u(x) = \Delta(x)\mathbf{1}\{\tilde{x}_p > 0\} - \Delta(x)\mathbf{1}\{\tilde{x}_p \leq 0\}$ (resp. $u(x) = -\Delta(x)\mathbf{1}\{\tilde{x}_p > 0\} + \Delta(x)\mathbf{1}\{\tilde{x}_p \leq 0\}$). Hence $E(\tilde{x}_p u(x))$ lies between $-E(|\tilde{x}_p| \Delta(x))$ and $E(|\tilde{x}_p| \Delta(x))$ and the difference $\beta_p - \beta_p^*$ varies in:

$$\left] -\frac{E(|\tilde{x}_p| \Delta(x))}{E(\tilde{x}_p^2)}, \frac{E(|\tilde{x}_p| \Delta(x))}{E(\tilde{x}_p^2)} \right].$$

To show the reciprocal, consider any β_p in

$$\left] \beta_p^* - \frac{E(|\tilde{x}_p| \Delta(x))}{E(\tilde{x}_p^2)}; \beta_p^* + \frac{E(|\tilde{x}_p| \Delta(x))}{E(\tilde{x}_p^2)} \right].$$

Denote

$$\lambda = \frac{E(\tilde{x}_p^2)}{E(|\tilde{x}_p| \Delta(x))} (\beta_p - \beta_p^*) \in]-1, 1[.$$

Consider $u(x) = \lambda \Delta(x)$ when $\tilde{x}_p > 0$ and $u(x) = -\lambda \Delta(x)$ otherwise which means that

$$\frac{E(\tilde{x}_p u(x))}{E(\tilde{x}_p^2)} = (\beta_p - \beta_p^*).$$

Function $u(x)$ takes its values in $]-\Delta(x), \Delta(x)[$ and therefore satisfies point (ii) of Theorem 2. Thus, there exists $\beta \in B$ such that its last component is β_p .

The adaptation to the general IV case uses the generalized transformation:

$$\begin{aligned}\tilde{x}_p &= z(E(z'z))^{-1}E(z'x_p) - \\ &\quad z(E(z'z))^{-1}E(z'x_{-p}) [E(x'_{-p}z)(E(z'z))^{-1}E(z'x_{-p})]^{-1} E(x'_{-p}z)(E(z'z))^{-1}E(z'x_p)\end{aligned}$$

Generally speaking, the estimation of B_p requires the estimation of $E(|\tilde{x}_p| \Delta(x, z))$. Given this fact, it is worth emphasizing that $\Delta(x, z)$ can be rewritten as $E(\tilde{y}_\Delta | x, z)$ where

$$\tilde{y}_\Delta = \frac{\mu_k \cdot y}{p_k(x, z)} + \frac{v_K - v_{K-1}}{2},$$

with

$$\mu_k = \frac{(v_k - v_{k-1} - (v_{k+1} - v_k))}{2} \text{ for } k = 2, \dots, K-1$$

and $\mu_1 = \mu_K = 0$. Specifically,

$$\begin{aligned}\Delta(x, z) &= \frac{1}{2} \sum_{k=2}^K [(v_k - v_{k-1})(G_k(x, z) - G_{k-1}(x, z))] \\ &= \frac{1}{2} [(v_2 - v_1)G_2(x, z) + (v_3 - v_2)(G_3(x, z) - G_2(x, z)) + \dots \\ &\quad \dots + (v_{K-1} - v_{K-2})(G_{K-1}(x, z) - G_{K-2}(x, z)) + (v_K - v_{K-1})(1 - G_{K-1}(x, z))] \\ &= \frac{1}{2} \sum_{k=2}^{K-1} (v_k - v_{k-1} - (v_{k+1} - v_k))G_k(x, z) + \frac{v_K - v_{K-1}}{2} \\ &= \frac{1}{2} \sum_{k=2}^{K-1} (v_k - v_{k-1} - (v_{k+1} - v_k))E(y | v = v_k, x, z) + \frac{v_K - v_{K-1}}{2} \\ &= E\left(\frac{\mu_k \cdot y}{p_k(x, z)} \mid x, z\right) + \frac{v_K - v_{K-1}}{2} = E(\tilde{y}_\Delta \mid x, z)\end{aligned}$$

Hence, $E(|\tilde{x}_p| \Delta(x, z))$ can be rewritten $E(|\tilde{x}_p| \tilde{y}_\Delta)$ which means that the estimation of the upper and lower bounds of B_p only requires [1] the construction of the transform \tilde{y}_Δ , [2] an estimation of the residual \tilde{x}_p and [3] the linear regression of \tilde{y}_Δ on $|\tilde{x}_p|$.

A.4 Proof of Lemma 5

First, by construction, $B \subset I(B)$.

Second, consider $\beta_0 \notin B$. By the separating hyperplane theorem (Takayama, 1974), the fact that B is a non empty closed and convex bounded set implies that there exists an hyperplane H_0 which contains β_0 and which intersection with B is empty ($H_0 \cap B = \emptyset$). Denote q_0 the orthogonal vector to H_0 so that $H_0 = \{\beta \in \mathbb{R}^p : q_0' \beta = q_0' \beta_0\}$. By construction,

$$\begin{aligned}H_0 \cap B &= \emptyset \implies H_0 \cap B_{q_0} = \emptyset \\ \implies \beta_0 &\notin I_{q_0}(B) \implies \beta_0 \notin \cap_{\|q\|=1} I_q(B) = I(B)\end{aligned}$$

Hence, $I(B) \subset B$. ■

B Proofs in Section 4

B.1 Proof of Theorem 7

Consider a vector of parameters β and a conditional probability distribution $\Pr(y = 1 \mid v^*, x, z)$ (denoted $G_{v^*}(x, z)$) which is non-decreasing in v^* .

(*Necessity*) We prove that (i) implies (ii). Denote, $F_v(\cdot \mid x, z, v^*)$, and $F_\varepsilon(\cdot \mid x, z)$, two conditional distribution functions satisfying (i). By Assumption $R(vi)$, $F_v(\cdot \mid x, z, v^*)$ is absolutely continuous and its density function is denoted f_v . By assumption (i), $(\beta, F_\varepsilon(\cdot \mid x, z))$ satisfies condition $(L1^*)$, $(L2)$ and $(L3)$ and $\{G_k(x, z)\}_{k=1, \dots, K-1}$ is its image through transformation (LV) .

For the sake of clarity, set $w = -(x\beta + \varepsilon)$ so that $y = \mathbf{1}\{v > w\}$ and the support of w is a subset of $[v_1, v_K[$ by $(L.2)$. The variable w is conditionally (on (x, z)) independent of v and v^* and the corresponding conditional distribution is:

$$F_w(w \mid x, z) = 1 - F_\varepsilon(-(x\beta + w) \mid x, z)$$

The conditional probability of occurrence of $y = 1$ in the k -th interval ($v^* = k$ in $\{1, \dots, K - 1\}$) is,

$$G_k(x, z) = \int_{v_k}^{v_{k+1}} E(\mathbf{1}\{v > w \mid v, v^* = k, x, z\} f_v(v \mid k, x, z) dv$$

which yields the convolution equation:

$$G_k(x, z) = \int_{v_k}^{v_{k+1}} F_w(v \mid x, z) f_v(v \mid k, x, z) dv. \quad (\text{B.1})$$

Note that this condition implies:

$$F_w(v_k \mid x, z) \leq G_k(x, z) \leq F_w(v_{k+1} \mid x, z). \quad (\text{B.2})$$

with a strict inequality on the right if $F_w(v_k \mid x, z) < F_w(v_{k+1} \mid x, z)$ because F_v is absolutely continuous and F_w is continuous on the right (CADLAG).

To prove (5), write $E(\bar{y} \mid x, z)$ as

$$\sum_{v^*=1, \dots, K-1} \int_{\Omega(v \mid v^*, x, z)} \int_{\Omega(w \mid v^*, v, x, z)} [\bar{y} \cdot p_{v^*}(x, z) \cdot f_v(v \mid v^*, x, z) dv dF_w(w \mid v^*, v, x, z)].$$

Using the definition of \bar{y} , the term $p_{v^*}(x, z)$ cancels out and using condition $(L.1^*)$, the integral over dw on the one hand, and the sum and other integral on the other hand, can be permuted:

$$\int_{\Omega(w \mid x, z)} \left[\sum_{v^* \in \{1, \dots, K-1\}} \delta(v^*) \int_{\Omega(v \mid v^*, x, z)} \mathbf{1}(v > w) f_v(v \mid v^*, x, z) dv \right] dF_w(w \mid x, z) - v_K. \quad (\text{B.3})$$

Evaluate first the inner integral with respect to v . As the support of w is included in $[v_1, v_K[$, we can define for any value of w in its support, an integer function $j(w)$ in $\{1, \dots, K - 1\}$,

such that $v_{j(w)} \leq w < v_{j(w)+1}$. Distinguish three cases. First, when $v^* < j(w)$, the whole conditional support of v lies below w and,

$$\int_{\Omega_{(v|v^*,x,z)}} \mathbf{1}(v > w) f_v(v | v^*, x, z) dv = 0.$$

while when $v^* > j(w)$, the whole conditional support of v lies strictly above w and thus:

$$\int_{\Omega_{(v|v^*,x,z)}} \mathbf{1}(v > w) f_v(v | v^*, x, z) dv = 1.$$

Last when $v^* = j(w)$,

$$\int_{\Omega_{(v|v^*,x,z)}} \mathbf{1}(v > w) f_v(v | v^*, x, z) dv = 1 - F_v(w | v^*, x, z).$$

Summing over values of v^* ,

$$\begin{aligned} \sum_{v^* \in \{1, \dots, K-1\}} \delta(v^*) \int_{\Omega_{(v|v^*,x,z)}} \mathbf{1}(v > w) f_v(v | v^*, x, z) dv \\ = -F_v(w | v_{j(w)}, x, z)(v_{j(w)+1} - v_{j(w)}) + v_K - v_{j(w)}. \end{aligned}$$

Replacing in (B.3) and integrating w.r.t. w , implies that:

$$E(\bar{y} | x, z) = -E(w | x, z) - u^*(x, z) = x\beta + E(\epsilon | x, z) - u^*(x, z). \quad (\text{B.4})$$

where

$$u^*(x, z) = \int_{\Omega_{(w|x,z)}} (F_v(w | v_{j(w)}, x, z)(v_{j(w)+1} - v_{j(w)}) + v_{j(w)} - w) dF_w(w | x, z).$$

Integrating (B.4) with respect to x, z and using condition (L.3) yields condition (5).

To finish the proof, upper and lower bounds for $u^*(x, z)$ are now provided. Let write,

$$u^*(x, z) = \sum_{k=1}^{K-1} (v_{k+1} - v_k) \phi_k(x, z) \quad (\text{B.5})$$

where:

$$\phi_k(x, z) = \int_{v_k}^{v_{k+1}} (F_v(w | k, x, z) + \frac{v_k - w}{v_{k+1} - v_k}) dF_w(w | x, z). \quad (\text{B.6})$$

By integration by parts, the first term is:

$$\phi_k(x, z) = \int_{v_k}^{v_{k+1}} \left(\frac{1}{v_{k+1} - v_k} - f_v(w | k, x, z) \right) F_w(w | x, z) dw$$

Therefore, using the convolution equation (B.1),

$$\phi_k(x, z) = -G_k(x, z) + \int_{v_k}^{v_{k+1}} \frac{F_w(w | x, z)}{v_{k+1} - v_k} dw.$$

Using (B.2) implies

$$G_{k-1}(x, z) - G_k(x, z) \leq \phi_k(x, z) \leq G_{k+1}(x, z) - G_k(x, z).$$

where at least one inequality on the right and one inequality on the left are strict since there exists at least one k such that $F_w(w = -(x\beta + v_{k+1}) | x, z) - F_w(w = -(x\beta + v_k) | x, z) > 0$. Therefore:

$$\underline{\Delta}^*(x, z) < u^*(x, z) < \overline{\Delta}^*(x, z).$$

where the definitions of $\overline{\Delta}^*(x, z)$ and $\underline{\Delta}^*(x, z)$ correspond to those given in the body of the Theorem.

(*Sufficiency*) We now prove that (ii) implies (i). Denote $u^*(x, z)$ in $]\underline{\Delta}^*(x, z), \overline{\Delta}^*(x, z)[$ such that

$$E(z'(x\beta - \bar{y})) = E(z'u^*(x, z))$$

We are going to prove that there exists a distribution function of $w = -(x\beta + \varepsilon)$ and a distribution function of v such that $(\beta, F_\varepsilon(\cdot | x, z))$ satisfies (L.1*, L.2, L.3) and $G_{v^*}(x, z)$ is the image of $(\beta, F_\varepsilon(\cdot | x, z))$ through the transformation (LV).

To begin with, we are going to construct w . We proceed in three steps.

First, we choose a sequence of functions $H_k(x, z)$ such that $H_1 = 0$, $H_K = 1$, and such that:

$$H_k(x, z) \leq G_k(x, z) \leq H_{k+1}(x, z), \text{ for } k \in \{1, \dots, K-1\} \quad (\text{B.7})$$

where at least one inequality on the right is strict and:

$$\sum_{k=1}^{K-1} (v_{k+1} - v_k)(H_k(x, z) - G_k(x, z)) < u^*(x, z) < \sum_{k=1}^{K-1} (v_{k+1} - v_k)(H_{k+1}(x, z) - G_k(x, z))$$

Consider for instance

$$\theta(x, z) > \frac{u^*(x, z)}{\underline{\Delta}^*(x, z)}, 1 - \theta(x, z) > \frac{u^*(x, z)}{\overline{\Delta}^*(x, z)},$$

for instance, $\theta(x, z) = \frac{\overline{\Delta}^*(x, z) - u^*(x, z)}{\overline{\Delta}^*(x, z) - \underline{\Delta}^*(x, z)}$. By construction $\theta(x, z) \in]0, 1]$ and one checks that

$$H_k(x, z) = \theta(x, z)G_{k-1}(x, z) + (1 - \theta(x, z))G_k(x, z)$$

satisfies the two previous conditions. Generally speaking, the closer $u^*(x, z)$ is from the lower bound $\underline{\Delta}^*(x, z)$, the closer is H_k to G_{k-1} , and the closer $u^*(x, z)$ is from the upper bound $\overline{\Delta}^*(x, z)$, the closer is H_k to G_k .

Secondly, we consider κ a discrete random variable which support is $\{1, \dots, K-1\}$, which is independent of v^* (a.e. $F_{x,z}$) and which conditional on (x, z) distribution is:

$$Pr(\kappa = k | x, z) = H_{k+1}(x, z) - H_k(x, z). \quad (\text{B.8})$$

Thirdly, we consider λ a random variable which support is $]0, 1[$, which is independent of v^* (a.e. $F_{x,z}$) and which conditional (on (x, z)) expectation is:

$$E(\lambda \mid x, z) = \frac{\sum_{k=1}^{K-1} (v_{k+1} - v_k)(H_{k+1}(x, z) - G_k(x, z)) - u^*(x, z)}{\sum_{k=1}^{K-1} (v_{k+1} - v_k)(H_{k+1}(x, z) - H_k(x, z))} \quad (\text{B.9})$$

For instance, λ can be chosen discrete with a mass point on

$$\lambda_0(x, z) = \frac{\sum_{k=1}^{K-1} (v_{k+1} - v_k)(H_{k+1}(x, z) - G_k(x, z)) - u^*(x, z)}{\sum_{k=1}^{K-1} (v_{k+1} - v_k)(H_{k+1}(x, z) - H_k(x, z))}.$$

Given the constraints on the $H_k(x, z)$ and given that $u^*(x, z)$ is in in $]\underline{\Delta}^*(x, z), \overline{\Delta}^*(x, z)[$, $\lambda_0(x, z)$ belongs to $]0, 1[$.

Within this framework, we can define w as:

$$w = (1 - \lambda)v_\kappa + \lambda v_{\kappa+1}$$

By construction, the support of w is $[v_1, v_K[$ and w is independent of v^* conditionally on (x, z) because both λ and κ are. Hence, $\varepsilon = -(x\beta + w)$ satisfies (L.1) and (L.2).

To construct v , we first introduce a random variable η which support is $[0, 1[$, which is absolutely continuous, which is defined conditionally on (k, x, z) , which is independent of λ and such that:

$$\int_0^1 F_\lambda(\eta \mid x, z) \cdot f_\eta(\eta \mid k, x, z) d\eta = \frac{G_k(x, z) - H_k(x, z)}{H_{k+1}(x, z) - H_k(x, z)} \in [0, 1[$$

where $F_\lambda(\cdot \mid x, z)$ denotes the distribution of λ conditional on (x, z) .

For instance, when λ is chosen discrete with a mass point on $\lambda_0(x, z)$, we simply have to chose η such that

$$F_\eta(\lambda_0(x, z) \mid x, z) = \frac{H_{k+1}(x, z) - G_k(x, z)}{H_{k+1}(x, z) - H_k(x, z)}.$$

Within this framework, we define v by the following expression:

$$v = v_k + (v_{k+1} - v_k)\eta$$

Having defined w and v , we are now going to prove that the image of $(\beta, F_w(\cdot \mid x, z))$ through (LV) is $G_{v^*}(x, z)$ because it satisfies equation (B.1):

$$\begin{aligned} & \int_{v_k}^{v_{k+1}} F_w(v \mid x, z) \cdot f_v(v \mid k, x, z) dv = H_k(x, z) + \\ & + (H_{k+1}(x, z) - H_k(x, z)) \int_{v_k}^{v_{k+1}} \Pr(w = (1 - \lambda)v_k + \lambda v_{k+1} \leq v \mid x, z) \cdot f_v(v \mid k, x, z) dv = \\ & H_k(x, z) + (H_{k+1}(x, z) - H_k(x, z)) \int_0^1 \Pr(\lambda \leq \eta \mid x, z) \cdot f_\eta(\eta \mid k, x, z) d\eta = G_k(x, z) \end{aligned}$$

The last condition to prove is (L.3). Rewrite equation (B.6), for almost any (x, z) ,

$$\begin{aligned}\phi_k(x, z) &= -G_k(x, z) + \int_{v_k}^{v_{k+1}} \frac{F_w(w | x, z)}{v_{k+1} - v_k} dw \\ &= -G_k(x, z) + H_{k+1}(x, z) - (H_{k+1}(x, z) - H_k(x, z))E(\lambda | x, z).\end{aligned}$$

Therefore,

$$\begin{aligned}\sum_{k=1}^{K-1} (v_{k+1} - v_k) \phi_k(x, z) &= \sum_{k=1}^{K-1} (v_{k+1} - v_k) (H_{k+1}(x, z) - G_k(x, z)) \\ &\quad - \sum_{k=1}^{K-1} (v_{k+1} - v_k) (H_{k+1}(x, z) - H_k(x, z)) E(\lambda | x, z) = u^*(x, z).\end{aligned}$$

using equation (B.9). Plugging (5) in (B.4) yields $E(z'\varepsilon) = 0$ that is (L.3).

B.2 Proof of Theorem 8

We use large parts of the proof of Theorem 7:

(*Necessity*) Same as the proof of Theorem 7 until equation (B.6) that we rewrite as:

$$\phi_k(x, z) = \int_{v_k}^{v_{k+1}} \left(\Phi(v | k, x, z) - \frac{v - v_k}{v_{k+1} - v_k} \right) dF_w(v | x, z).$$

We then have first:

$$\begin{aligned}\phi_k(x, z) &= \int_{v_k}^{v_{k+1}} \left(1 - \frac{\frac{v - v_k}{v_{k+1} - v_k}}{\Phi(v | k, x, z)} \right) \Phi(v | k, x, z) dF_w(v | x, z) \\ &\leq \sup_{v \in]v_k, v_{k+1}[} \left(1 - \frac{\frac{v - v_k}{v_{k+1} - v_k}}{\Phi(v | k, x, z)} \right) \int_{v_k}^{v_{k+1}} \Phi(v | k, x, z) dF_w(v | x, z) \\ &= \xi_k^U(x, z) \int_{v_k}^{v_{k+1}} \Phi(v | k, x, z) dF_w(v | x, z).\end{aligned}$$

where $\xi_k^U(x, z)$ is defined in the text. Equation (B.1) delivers:

$$\begin{aligned}&\int_{v_k}^{v_{k+1}} \Phi(v | k, x, z) dF_w(v | x, z). \\ &= \int_{v_k}^{v_{k+1}} d[\Phi(v | k, x, z) F_w(v | x, z)] - \int_{v_k}^{v_{k+1}} F_w(v | x, z) d\Phi(v | k, x, z) \\ &= \int_{v_k}^{v_{k+1}} d[\Phi(v | k, x, z) F_w(v | x, z)] - G_k(x, z).\end{aligned}$$

Hence, using $F_w(v_{k+1} | x, z) \leq G_{k+1}(x, z)$, we have,

$$\phi_k(x, z) \leq \xi_k^U(x, z) \cdot (G_{k+1}(x, z) - G_k(x, z)).$$

The derivation of the lower bound follows the same logic:

$$\begin{aligned}\phi_k(x, z) &\geq \inf_{v \in]v_k, v_{k+1}[} \left(-1 - \frac{\frac{v-v_{k+1}}{v_{k+1}-v_k}}{1 - \Phi(v | k, x, z)}\right) \int_{v_k}^{v_{k+1}} (1 - \Phi(v | k, x, z)) dF_w(v | x, z) \\ &\geq \xi_k^L(x, z) \left[\int_{v_k}^{v_{k+1}} d[(1 - \Phi(v | k, x, z)) F_w(v | x, z)] + G_k(x, z) \right]\end{aligned}$$

where $\xi_k^L(x, z)$ is defined in the text. Hence, using $F_w(v_k | x, z) \geq G_k(x, z)$, we have

$$\phi_k(x, z) \geq \xi_k^L(x, z)(G_k(x, z) - G_{k-1}(x, z)).$$

Therefore, using the definition of $u^*(x, z)$ (B.5), we have:

$$\underline{\Delta}_{\Phi}^*(x, z) \leq u^*(x, z) \leq \overline{\Delta}_{\Phi}^*(x, z) \quad (\text{B.10})$$

where $\underline{\Delta}_{\Phi}^*(x, z)$ and $\overline{\Delta}_{\Phi}^*(x, z)$ are defined in the text.

(*Sufficiency*) We now prove that (ii) implies (i). We assume that there exists $u^*(x, z)$ in $[\underline{\Delta}_{\Phi}^*(x, z), \overline{\Delta}_{\Phi}^*(x, z)]$ such that

$$E(z'(x\beta - \bar{y})) = E(z'u^*(x, z))$$

Under this assumption, we are going to prove that there exists a distribution function of the random term ε such that $(\beta, F_{\varepsilon}(\cdot | x, z))$ satisfies (L.1*, L.2, L.3) and $G_{v^*}(x, z)$ is the image of $(\beta, F_{\varepsilon}(\cdot | x, z))$ through the transformation (LV), when the distribution function of the special regressor v is $\Phi(v | k, x, z)$. As in the proof of Theorem 7, we proceed by construction in three steps.

First, choose a sequence of functions $H_k(x, z)$ such that $H_1 = 0$, $H_K = 1$, and for any k in $\{1, \dots, K-1\}$ such as:

$$H_k(x, z) \leq G_k(x, z) < H_{k+1}(x, z). \quad (\text{B.11})$$

and such as:

$$\begin{aligned}&\sum_{k=1}^{K-1} (v_{k+1} - v_k) \xi_k^L(x, z) (G_k(x, z) - H_k(x, z)) \leq u^*(x, z) \\ &\leq \sum_{k=1}^{K-1} (v_{k+1} - v_k) \xi_k^U(x, z) (H_{k+1}(x, z) - G_k(x, z))\end{aligned}$$

If $\xi_k^L(x, z) < 0$ and $\xi_k^U(x, z) > 0$, the closer $u^*(x, z)$ is from the lower bound $\underline{\Delta}_{\Phi}^*(x, z)$, the closer is H_k to G_{k-1} , and the closer $u^*(x, z)$ is from the upper bound $\overline{\Delta}_{\Phi}^*(x, z)$, the closer is H_k to G_k .

Decompose now $u^*(x, z)$ into $\phi_k^*(x, z)$ such that:

$$u^*(x, z) = \sum_{k=1}^{K-1} (v_{k+1} - v_k) \phi_k^*(x, z)$$

and such that the bounds on u^* can be translated into:

$$\xi_k^L(x, z)(G_k(x, z) - H_k(x, z)) \leq \phi_k^*(x, z) \leq \xi_k^U(x, z)(H_{k+1}(x, z) - G_k(x, z)) \quad (\text{B.12})$$

There are many decompositions of this type. Choose one.

Second, consider κ a discrete random variable which support is $\{1, \dots, K - 1\}$, which is independent of v^* (a.e. $F_{x,z}$) and which conditional on (x, z) distribution is:

$$Pr(\kappa = k \mid x, z) = H_{k+1}(x, z) - H_k(x, z). \quad (\text{B.13})$$

Consider also $K - 1$ random variable λ_k which support is $]0, 1[$, which are independent of v^* (a.e. $F_{x,z}$) and which conditional (on (x, z)) expectation is:

$$E(\lambda_k \mid x, z) = \frac{H_{k+1}(x, z) - G_k(x, z) - \phi_k^*(x, z)}{H_{k+1}(x, z) - H_k(x, z)} \quad (\text{B.14})$$

and such that:

$$\int_0^1 (\Phi_v(\lambda v_k + (1 - \lambda)v_{k+1} \mid k, x, z) - \frac{v - v_k}{v_{k+1} - v_k}) dF_{\lambda_k}(\lambda \mid x, z) = \frac{\phi_k^*(x, z)}{H_{k+1}(x, z) - H_k(x, z)}$$

Given constraints (B.11) and (B.12), it is always possible to construct such a random variable.

Finally, define the random variable:

$$w = (1 - \lambda)v_\kappa + \lambda v_{\kappa+1}$$

By construction, the support of w is $[v_1, v_K[$ and w is independent of v^* conditionally on (x, z) because all λ_k s and κ are. Hence, $\varepsilon = -(x\beta + w)$ satisfies (L.1) and (L.2).

Finish the proof as in Theorem 7.

B.3 Proof of Corollary 9

(*Necessity*) Let the conditional distribution of v , Φ_0 , be piece-wise uniform by intervals, $v^* = k$. Then, for any $k = 1, \dots, K - 1$, $\xi_k^U(x, z) = \xi_k^L(x, z) = 0$. Using Theorem 8 yields that $\underline{\Delta}_\Phi^*(x, z) = \overline{\Delta}_\Phi^*(x, z) = 0$ and therefore $u^*(x, z) = 0$. Identification of β is exact and its value is given by the moment condition (5).

(*Sufficiency*) By contraposition; Assume that there exists $k \in \{1, \dots, K - 1\}$, a measurable set A included in $[v_k, v_{k+1}[$ with positive Lebesgue measure and a measurable set S of elements (x, z) with positive probability $F_{x,z}(S) > 0$ such that $\Phi(v \mid k, x, z)$ is different from a uniform distribution function on A for any (x, z) in S . Because Φ is absolutely continuous (*ID(ii)*), and for the sake of simplicity assume that:

$$\forall v \in A; \forall (x, z) \in S; \Phi(v \mid k, x, z) - \frac{v - v_k}{v_{k+1} - v_k} > 0$$

Because $\xi_k^U(x, z) > 0$, we can always construct a function $u_1^*(x, z)$ which is strictly positive on S satisfying the conditions of Theorem 8. Thus $E(z'u_1^*(x, z)) \neq 0$ and the moment condition

(5) can be used to construct parameter β_1 . It implies that the identification set B contains at least two different parameters β , i.e. the one corresponding to $u^*(x, z) = 0$ and the one corresponding to $u_1^*(x, z)$ (and in fact the whole real line between them as B is convex).

Interpretation:

Consider a observable variable v_0 drawn conditionally on v^* in a uniform distribution in $[v_k, v_{k+1}]$. Write an auxiliary model as:

$$y = \mathbf{1}\{v_0 + x\beta + \varepsilon_0 > 0\}$$

where by construction:

$$\varepsilon_0 = \varepsilon + v - v_0.$$

Note first that v and v_0 are independent conditional on (v^*, x, z) . Second, that the auxiliary model now is a binary model with a continuous special regressor. Third, that the discrete-type transformation \bar{y} of the data is equal up to a constant term to the continuous-type transformation of the data *i.e.*:

$$\bar{y} = \frac{v_{k+1} - v_k}{p_{v^*}(x, z)} y - v_K = \frac{y}{f_{v_0}(v_0, v_*, x, z)} - v_K = \tilde{y} + cst$$

since by construction:

$$f_{v_0}(v_0, v_*, x, z) = \frac{p_{v^*}(x, z)}{v_{k+1} - v_k}$$

The method of Lewbel (2000) can be applied to the auxiliary model and data (y, v_0, v^*, x, z) to get consistent estimates of parameter β if several conditions hold. We shall only check the first of these conditions which is partial independence. What should hold is:

$$F(\varepsilon_0 | v_0, v^*, x, z) = F(\varepsilon_0 | v^*, x, z)$$

For convenience, omit the conditioning on (v^*, x, z) . Thus:

$$\begin{aligned} F(\varepsilon_0 | v_0) &= \Pr(\varepsilon + v - v_0 \leq \varepsilon_0 | v_0) \\ &= \int f_\varepsilon(\varepsilon | v_0) f_v(\varepsilon_0 - (\varepsilon - v_0) | v_0) d\varepsilon \end{aligned}$$

As v_0 is a random draw $f_\varepsilon(\varepsilon | v_0) = f_\varepsilon(\varepsilon)$ and $f_v(v | v_0) = f_v(v)$, we have:

$$F(\varepsilon_0 | v_0) = \int f_\varepsilon(\varepsilon) f_v(\varepsilon_0 - (\varepsilon - v_0)) d\varepsilon$$

The only dependence on v_0 occurs through the density function of v and it is in the case of a uniform distribution only that partial independence holds:

$$F(\varepsilon_0 | v_0) = F(\varepsilon_0).$$

The other conditions should be checked and this is the large support one which “creates” the bias in the intercept term.

B.4 Proof of Corollary 12

Same as Corollary 3 except that the maximisation of $E(\tilde{x}_p u^*(x, z))$ is obtained when:

$$u^*(x, z) = \mathbf{1}\{\tilde{x}_p \leq 0\} \underline{\Delta}^*(x, z) + \mathbf{1}\{\tilde{x}_p > 0\} \overline{\Delta}^*(x, z)$$

and the minimization of such an expression is obtained when:

$$u^*(x, z) = \mathbf{1}\{\tilde{x}_p > 0\} \underline{\Delta}^*(x, z) + \mathbf{1}\{\tilde{x}_p \leq 0\} \overline{\Delta}^*(x, z)$$

Furthermore, we have:

$$\begin{aligned} \overline{\Delta}^*(x, z) &= \sum_{k=1}^{K-1} [(v_{k+1} - v_k)(G_{k+1}(x, z) - G_k(x, z))] \\ &= [(v_2 - v_1)(G_2(x, z) - G_1(x, z)) + (v_3 - v_2)(G_3(x, z) - G_2(x, z)) + \dots \\ &\quad \dots + (v_{K-1} - v_{K-2})(G_{K-1}(x, z) - G_{K-2}(x, z)) + (v_K - v_{K-1})(1 - G_{K-1}(x, z))] \\ &= -(v_2 - v_1)G_1(x, z) + \sum_{k=2}^{K-1} (v_k - v_{k-1} - (v_{k+1} - v_k))G_k(x, z) + v_K - v_{K-1} \\ &= \sum_{k=1}^{K-1} (v_k - v_{k-1} - (v_{k+1} - v_k))E(y \mid v = v_k, x, z) + v_K - v_{K-1} \\ &= E\left(\frac{\theta_{U,k} \cdot y}{p_k(x, z)} \mid x, z\right) + v_K - v_{K-1} = E(\bar{y}_U \mid x, z) \end{aligned}$$

where by convention $v_0 = v_1$. Similarly:

$$\begin{aligned} \underline{\Delta}^*(x, z) &= \sum_{k=1}^{K-1} [(v_{k+1} - v_k)(G_{k-1}(x, z) - G_k(x, z))] \\ &= [-(v_2 - v_1)G_1(x, z) + (v_3 - v_2)(G_1(x, z) - G_2(x, z)) + \dots \\ &\quad \dots + (v_{K-1} - v_{K-2})(G_{K-3}(x, z) - G_{K-2}(x, z)) + (v_K - v_{K-1})(G_{K-2}(x, z) - G_{K-1}(x, z))] \\ &= \sum_{k=1}^{K-2} (v_{k+2} - v_{k+1} - (v_{k+1} - v_k))G_k(x, z) - (v_K - v_{K-1})G_{K-1}(x, z) \\ &= \sum_{k=1}^{K-1} (v_{k+2} - v_{k+1} - (v_{k+1} - v_k))E(y \mid v = v_k, x, z) \\ &= E\left(\frac{\theta_{L,k} \cdot y}{p_k(x, z)} \mid x, z\right) = E(\bar{y}_L \mid x, z) \end{aligned}$$

if the convention $v_{K+1} = v_K$ is adopted.




Figure 1: A graphical argument for set-identification

Table 1: Simple experiment, Interval Data: Sensitivity to Bandwidth

Lower and upper estimated bounds with standard errors

Error Decomposition: Decentering, Adjusted Length and Sampling Error

Nobs	Bwidth	Intercept				Variable			
		Dec	AL	ASE	RMSEI	Dec	AL	ASE	RMSEI
100	1.000	0.554	0.396	0.398	0.789	-0.041	0.353	0.491	0.606
	1.500	0.410	0.404	0.325	0.661	-0.068	0.352	0.411	0.545
	3.000	0.247	0.401	0.266	0.541	-0.079	0.337	0.327	0.476
	5.000	0.182	0.383	0.281	0.508	-0.090	0.321	0.322	0.464
200	1.000	0.358	0.428	0.204	0.594	0.007	0.375	0.267	0.461
	1.500	0.218	0.422	0.172	0.505	-0.039	0.367	0.225	0.432
	3.000	0.079	0.412	0.144	0.444	-0.101	0.350	0.191	0.412
	5.000	0.046	0.410	0.140	0.435	-0.126	0.347	0.202	0.420
500	1.000	0.096	0.411	0.094	0.433	-0.053	0.357	0.122	0.381
	1.500	0.025	0.406	0.085	0.415	-0.088	0.349	0.106	0.375
	3.000	-0.038	0.400	0.080	0.410	-0.129	0.340	0.104	0.378
	5.000	-0.050	0.399	0.080	0.410	-0.138	0.338	0.113	0.383
1000	1.000	-0.006	0.402	0.059	0.406	-0.094	0.345	0.073	0.364
	1.500	-0.044	0.399	0.056	0.405	-0.119	0.339	0.067	0.365
	3.000	-0.076	0.396	0.055	0.407	-0.146	0.334	0.071	0.372
	5.000	-0.082	0.396	0.055	0.408	-0.151	0.334	0.078	0.374

Notes: The number of interval values is equal to 10. The simple experiment refers to the case where $\alpha = \rho = 0$. All details are reported in the text. Experimental results are based on 1000 replications. **LB** and **UB** refer to the estimated lower and upper bounds of intervals with their standard errors (**SE**). **Bwidth** refers to the bandwidth that is used. **Dec** stands for decentering of the mid-point of the interval that is, $(\mathbf{UB} + \mathbf{LB})/2$. **AL** is the adjusted length of the interval, $(\mathbf{UB} - \mathbf{LB})/2\sqrt{3}$. **ASE** is the sampling variability of bounds as defined in the text. The identity $\mathbf{Dec}^2 + \mathbf{AL}^2 + \mathbf{ASE}^2 = \mathbf{RMSEI}^2$ is shown in the text. **RMSEI** is the root mean square error integrated over the identified interval.

Table 2: Sensitivity to Normality, Interval Data

Lower and upper estimated bounds with standard errors

Nobs	Alpha	Intercept				Variable			
		LB	SE	UB	SE	LB	SE	UB	SE
100	0.000	-0.641	0.244	0.916	0.203	-0.820	0.502	0.605	0.314
100	0.333	-0.674	0.232	0.602	0.306	-0.697	0.285	0.351	0.238
100	0.667	-0.659	0.165	0.563	0.352	-0.403	0.351	0.569	0.337
100	1.000	-0.673	0.238	0.758	0.306	-0.136	0.483	1.077	0.332
200	0.000	-0.666	0.120	0.805	0.084	-0.695	0.162	0.582	0.152
200	0.333	-0.720	0.087	0.799	0.104	-0.639	0.199	0.664	0.131
200	0.667	-0.688	0.184	0.800	0.212	-0.502	0.194	0.730	0.153
200	1.000	-0.797	0.183	0.776	0.264	-0.160	0.165	1.106	0.161
500	0.000	-0.714	0.032	0.660	0.098	-0.728	0.133	0.460	0.139
500	0.333	-0.819	0.084	0.618	0.083	-0.650	0.134	0.614	0.087
500	0.667	-0.864	0.094	0.634	0.106	-0.552	0.066	0.747	0.096
500	1.000	-0.891	0.097	0.663	0.093	-0.486	0.165	0.889	0.162
1000	0.000	-0.790	0.039	0.598	0.032	-0.742	0.087	0.460	0.070
1000	0.333	-0.816	0.033	0.602	0.047	-0.635	0.088	0.590	0.056
1000	0.667	-0.844	0.064	0.606	0.049	-0.484	0.086	0.771	0.071
1000	1.000	-0.937	0.042	0.597	0.039	-0.430	0.102	0.941	0.097

Error Decomposition: Decentering, Adjusted Length and Sampling Error

Nobs	Alpha	Intercept				Variable			
		Dec	AL	ASE	RMSEI	Dec	AL	ASE	RMSEI
100	0	0.137	0.449	0.220	0.519	-0.107	0.411	0.401	0.584
100	0.33	-0.036	0.368	0.243	0.443	-0.173	0.302	0.246	0.427
100	0.66	-0.048	0.353	0.219	0.418	0.083	0.281	0.328	0.440
100	1	0.043	0.413	0.238	0.479	0.470	0.350	0.404	0.712
200	0	0.069	0.424	0.100	0.442	-0.057	0.369	0.130	0.395
200	0.33	0.039	0.438	0.094	0.450	0.013	0.376	0.164	0.411
200	0.66	0.056	0.429	0.196	0.475	0.114	0.356	0.160	0.407
200	1	-0.011	0.454	0.223	0.506	0.473	0.365	0.161	0.619
500	0	-0.027	0.397	0.067	0.403	-0.134	0.343	0.130	0.391
500	0.33	-0.100	0.415	0.083	0.435	-0.018	0.365	0.110	0.381
500	0.66	-0.115	0.433	0.100	0.459	0.098	0.375	0.081	0.396
500	1	-0.114	0.448	0.094	0.472	0.202	0.397	0.163	0.474
1000	0	-0.096	0.401	0.036	0.414	-0.141	0.347	0.077	0.383
1000	0.33	-0.107	0.409	0.040	0.425	-0.022	0.354	0.071	0.361
1000	0.66	-0.119	0.419	0.056	0.439	0.144	0.362	0.078	0.398
1000	1	-0.170	0.443	0.039	0.476	0.255	0.396	0.098	0.481

Notes: See Table 1 for main comments. Specifics are: The bandwidth is equal to 3.0. The **Alpha** column refers to the increasing amount of non-normality.

Table 3: Sensitivity to Endogeneity, Interval Data

Lower and upper estimated bounds with standard errors

Nobs	Rho	Intercept				Variable			
		LB	SE	UB	SE	L B	SE	UB	SE
100	0.000	-0.447	0.248	0.941	0.316	-0.663	0.356	0.505	0.340
100	0.333	-0.589	0.441	0.245	0.484	-0.600	0.516	0.362	0.522
100	0.667	-0.602	0.434	0.235	0.479	-0.626	0.515	0.343	0.522
100	1.000	-0.625	0.429	0.214	0.475	-0.659	0.513	0.315	0.515
200	0.000	-0.634	0.147	0.793	0.150	-0.708	0.212	0.506	0.200
200	0.333	-0.583	0.168	0.649	0.254	-0.684	0.274	0.461	0.265
200	0.667	-0.592	0.168	0.640	0.250	-0.697	0.279	0.453	0.268
200	1.000	-0.604	0.168	0.630	0.248	-0.714	0.281	0.442	0.268
500	0.000	-0.731	0.084	0.656	0.081	-0.719	0.118	0.460	0.107
500	0.333	-0.726	0.085	0.657	0.085	-0.757	0.125	0.501	0.120
500	0.667	-0.734	0.086	0.648	0.085	-0.768	0.125	0.494	0.118
500	1.000	-0.744	0.086	0.635	0.086	-0.782	0.127	0.482	0.118
1000	0.000	-0.762	0.058	0.610	0.055	-0.725	0.079	0.433	0.075
1000	0.333	-0.762	0.058	0.610	0.055	-0.764	0.081	0.467	0.077
1000	0.667	-0.769	0.059	0.601	0.056	-0.774	0.082	0.461	0.078
1000	1.000	-0.781	0.059	0.589	0.056	-0.790	0.083	0.451	0.078

Error Decomposition: Decentering, Adjusted Length and Sampling Error

Nobs	Rho	Intercept				Variable			
		Dec	AL	ASE	RMSEI	Dec	AL	ASE	RMSEI
100	0.000	0.247	0.401	0.266	0.541	-0.079	0.337	0.327	0.476
100	0.333	-0.172	0.241	0.448	0.537	-0.119	0.278	0.501	0.585
100	0.667	-0.183	0.242	0.442	0.536	-0.141	0.280	0.500	0.590
100	1.000	-0.205	0.242	0.437	0.540	-0.172	0.281	0.495	0.594
200	0.000	0.079	0.412	0.144	0.444	-0.101	0.350	0.191	0.412
200	0.333	0.033	0.356	0.193	0.406	-0.112	0.331	0.249	0.429
200	0.667	0.024	0.355	0.190	0.404	-0.122	0.332	0.253	0.435
200	1.000	0.013	0.356	0.190	0.404	-0.136	0.334	0.254	0.441
500	0.000	-0.038	0.400	0.080	0.410	-0.129	0.340	0.104	0.378
500	0.333	-0.034	0.399	0.083	0.409	-0.128	0.363	0.113	0.401
500	0.667	-0.043	0.399	0.083	0.410	-0.137	0.364	0.113	0.405
500	1.000	-0.055	0.398	0.083	0.410	-0.150	0.365	0.113	0.410
1000	0.000	-0.076	0.396	0.055	0.407	-0.146	0.334	0.071	0.372
1000	0.333	-0.076	0.396	0.055	0.407	-0.148	0.356	0.073	0.392
1000	0.667	-0.084	0.395	0.056	0.408	-0.157	0.356	0.074	0.396
1000	1.000	-0.096	0.396	0.056	0.411	-0.170	0.358	0.075	0.403

Notes: See Table 1 for main comments. Specifics are: The bandwidth is equal to 3.0. The **Rho** column refers to the increasing amount of endogeneity.

Table 4: Sensitivity to the Number of Intervals, Interval Data

Lower and upper estimated bounds with standard errors

Nobs	Intervals	Intercept				Variable			
		LB	SE	UB	SE	LB	SE	UB	SE
100	5	-1.104	0.195	0.998	0.177	-1.150	0.274	0.710	0.242
100	20	-0.166	0.508	0.299	0.527	-0.278	0.456	0.133	0.449
100	40	-1.090	0.542	-0.843	0.518	-0.360	0.486	-0.114	0.502
100	80	-2.153	0.551	-2.020	0.525	-0.533	0.533	-0.379	0.565
200	5	-1.173	0.129	0.901	0.114	-1.140	0.178	0.676	0.155
200	20	0.025	0.252	0.640	0.305	-0.290	0.307	0.209	0.311
200	40	-0.148	0.378	0.074	0.371	-0.147	0.332	0.054	0.329
200	80	-1.198	0.387	-1.078	0.384	-0.300	0.331	-0.177	0.335
500	5	-1.206	0.078	0.847	0.065	-1.138	0.109	0.643	0.089
500	20	-0.203	0.100	0.530	0.110	-0.351	0.150	0.258	0.149
500	40	0.157	0.164	0.484	0.191	-0.142	0.210	0.117	0.206
500	80	0.003	0.246	0.124	0.248	-0.066	0.213	0.036	0.215
1000	5	-1.215	0.054	0.831	0.045	-1.139	0.076	0.629	0.064
1000	20	-0.315	0.062	0.401	0.064	-0.370	0.095	0.225	0.091
1000	40	0.013	0.077	0.398	0.084	-0.187	0.124	0.125	0.122
1000	80	0.225	0.133	0.385	0.142	-0.074	0.151	0.052	0.151

Error Decomposition: Decentering, Adjusted Length and Sampling Error

Nobs	Intervals	Intercept				Variable			
		Dec	AL	ASE	RMSEI	Dec	AL	ASE	RMSEI
100	5	-0.053	0.607	0.174	0.633	-0.220	0.537	0.225	0.622
100	20	0.066	0.134	0.506	0.528	-0.073	0.119	0.443	0.464
100	40	-0.967	0.071	0.511	1.096	-0.237	0.071	0.475	0.536
100	80	-2.087	0.038	0.499	2.146	-0.456	0.044	0.504	0.681
200	5	-0.136	0.599	0.115	0.625	-0.232	0.524	0.146	0.592
200	20	0.333	0.177	0.269	0.463	-0.041	0.144	0.301	0.336
200	40	-0.037	0.064	0.368	0.375	-0.047	0.058	0.325	0.333
200	80	-1.138	0.035	0.373	1.198	-0.239	0.035	0.320	0.401
500	5	-0.179	0.593	0.068	0.623	-0.247	0.514	0.087	0.577
500	20	0.164	0.212	0.103	0.286	-0.047	0.176	0.143	0.231
500	40	0.320	0.095	0.174	0.376	-0.012	0.075	0.205	0.219
500	80	0.063	0.035	0.245	0.255	-0.015	0.029	0.212	0.215
1000	5	-0.192	0.591	0.047	0.623	-0.255	0.511	0.061	0.574
1000	20	0.043	0.207	0.062	0.220	-0.073	0.172	0.090	0.207
1000	40	0.205	0.111	0.079	0.247	-0.031	0.090	0.121	0.154
1000	80	0.305	0.046	0.136	0.337	-0.011	0.036	0.150	0.155

Notes: See Table 1 for main comments. Specifics are: The bandwidth is equal to 3.0. The **Intervals** column refers to the number of intervals of v .